

INFINITESIMAL DERIVED TORELLI THEOREM FOR K3 SURFACES

EMANUELE MACRÌ AND PAOLO STELLARI

ABSTRACT. We prove that the first order deformations of two smooth projective K3 surfaces are derived equivalent under a Fourier–Mukai transform if and only if there exists a special isometry of the total cohomology groups of the surfaces which preserves the Mukai pairing, an infinitesimal weight-2 decomposition and the orientation of a positive 4-dimensional space. This generalizes the derived version of the Torelli Theorem. Along the way we show the compatibility of the actions on Hochschild homology and singular cohomology of any Fourier–Mukai functor.

1. INTRODUCTION

A great deal of geometric information is encoded in the lattice and Hodge structures defined on the cohomology groups of a K3 surface (i.e. a smooth, simply-connected, projective surface with trivial canonical bundle). Just two results making this plain are the classical Torelli Theorem (see [5, 12, 23]) and its more recent categorical version, the Derived Torelli Theorem.

This latter theorem, in the final form resulting from the combination of [19, 22] and [9], asserts that any equivalence between the derived categories of coherent sheaves of two K3 surfaces induces a Hodge isometry on their cohomology groups which preserves the orientation of some positive four-space. The reverse implication is also true and follows from a detailed analysis of the geometry of moduli spaces of stable sheaves on K3 surfaces.

Since the deformation theory of K3 surfaces is well understood, one can wonder if, in an appropriate setting, the Derived Torelli Theorem can be extended to (at least) first order deformations. More precisely, one can ask if the equivalences of the derived categories of first order deformations of K3 surfaces can be detected by the existence of isometries of some kind of deformed lattice and Hodge structures on the total cohomology groups.

To state our answer to this question, we need to sketch briefly the categorical setting and some additional structures on the total cohomologies of K3 surfaces (which will be extensively described in Sections 3.1 and 3.2 respectively).

For a smooth projective (complex) variety X , all the abelian categories which are first order deformations of the abelian category $\mathbf{Coh}(X)$ of coherent sheaves on X are parametrized by the second Hochschild cohomology group $\mathrm{HH}^2(X)$. In particular, for each element $v \in \mathrm{HH}^2(X)$, one can produce (see [26]) an abelian category $\mathbf{Coh}(X, v)$ which is the first order deformation of $\mathbf{Coh}(X)$ in the direction v . The kernels of the Fourier–Mukai functors between the derived categories $\mathrm{D}^b(X_1, v_1)$ and $\mathrm{D}^b(X_2, v_2)$ of $\mathbf{Coh}(X_1, v_1)$ and $\mathbf{Coh}(X_2, v_2)$ are perfect complexes (i.e. complexes locally quasi-isomorphic to a finite complex of locally free sheaves) in $\mathrm{D}_{\mathrm{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$. Roughly speaking, the operator J changes the sign of a component of v_1 .

If X is a K3 surface, the total cohomology group $H^*(X, \mathbb{Z})$ tensored by $\mathbb{Z}[\epsilon]/(\epsilon^2)$ inherits a pairing and, chosen $v \in \mathrm{HH}^2(X)$, a weight-2 decomposition (via the Hochschild–Kostant–Rosenberg isomorphism). Such a $\mathbb{Z}[\epsilon]/(\epsilon^2)$ -module, endowed with these additional structures, is called *infinitesimal Mukai lattice* and it is denoted by $\tilde{H}(X, v, \mathbb{Z})$ (see Definition 3.5). This lattice is closely related to the one defined in [10] for the case of twisted K3 surfaces. The analogy is actually quite

2000 *Mathematics Subject Classification.* 18E30, 14J28, 13D10.

Key words and phrases. K3 surfaces, derived categories, deformations.

precise, since a K3 surface twisted by an element in its Brauer group can be thought as a complete (not just formal) deformation in a gerby direction.

An isomorphism $g : \tilde{H}(X_1, v_1, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(X_2, v_2, \mathbb{Z})$ of $\mathbb{Z}[\epsilon]/(\epsilon^2)$ -modules is a Hodge isometry if it preserves the pairings and the weight-2 decompositions. The Hodge isometries we will consider are the effective and orientation preserving ones, i.e. those induced by isomorphisms $H^*(X_1, \mathbb{Z}) \xrightarrow{\sim} H^*(X_2, \mathbb{Z})$ preserving the Mukai pairings, the natural weight-2 Hodge structures and the orientation of the spaces generated by the four positive directions in $H^*(X_1, \mathbb{Z})$ and $H^*(X_2, \mathbb{Z})$ (see the beginning of Section 3.2).

With this notation in mind we can state the main result of this paper, which is an infinitesimal version of the Derived Torelli Theorem.

Theorem 1.1. *Let X_1 and X_2 be smooth complex projective K3 surfaces and let $v_i \in \mathrm{HH}^2(X_i)$, with $i = 1, 2$. Then the following are equivalent:*

- (i) *There exists a Fourier–Mukai equivalence*

$$\Phi_{\tilde{\mathcal{E}}} : \mathrm{D}^b(X_1, v_1) \xrightarrow{\sim} \mathrm{D}^b(X_2, v_2)$$

with $\tilde{\mathcal{E}} \in \mathrm{D}_{\mathrm{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$.

- (ii) *There exists an orientation preserving effective Hodge isometry*

$$g : \tilde{H}(X_1, v_1, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(X_2, v_2, \mathbb{Z}).$$

As in the classical case, this result can be reformulated in a neat way for autoequivalences of Fourier–Mukai type. Indeed, Theorem 1.1 yields the existence of a surjective group homomorphism

$$\mathrm{Aut}^{\mathrm{FM}}(\mathrm{D}^b(X, v)) \longrightarrow \mathrm{O}_+(\tilde{H}(X, v, \mathbb{Z})),$$

where $\mathrm{Aut}^{\mathrm{FM}}(\mathrm{D}^b(X, v))$ is the group of equivalences of Fourier–Mukai type of the triangulated category $\mathrm{D}^b(X, v)$ and $\mathrm{O}_+(\tilde{H}(X, v, \mathbb{Z}))$ is the group of orientation preserving effective Hodge isometries. In the non-deformed case, Bridgeland gave in [2] a very nice conjectural description of the kernel of the previous morphism. As we will observe in Section 3.3, the same applies to the infinitesimal case.

The final goal of our investigation would be to generalize Theorem 1.1 to deformations of K3 surfaces of any order and possibly formal. A key step in this direction would be to show that one can deform to any order the abelian categories of coherent sheaves and, compatibly, the kernel of any Fourier–Mukai equivalence. There are already examples in the literature of deformations in this broader generality. This is the case of the Poincaré sheaf for an abelian variety and its dual (see [1]). Unfortunately, the argument there seems to be quite ad hoc and we cannot hope to apply those techniques to the situation we want to treat. A more general attempt to deal with this problem for abelian varieties has been pursued by D. Arinkin. Deformations of kernels of Fourier–Mukai equivalences were also studied in [9] for very special analytic directions in the case of K3 surfaces. For first order deformations, such a theory, which will be used in this paper, has been completely carried out in [26] (see Theorem 3.4).

Once this goal is achieved, one should be able to define a deformation functor DF_1 , over local Artin algebras, of families of derived categories of non-commutative and gerby K3 surfaces. The definition of DF_1 , for first-order-deformations, is the content of [26].

On the other hand, there exists a second deformation functor DF_2 of variations of weight-2 Hodge structures of the Mukai lattice, endowed with the Mukai pairing. This is because, there is a global versal analytic deformation, which is local analytically mini-versal, provided by a period domain. Once an appropriate definition of DF_1 is given, one should also have a morphism of functors $p : \mathrm{DF}_1 \rightarrow \mathrm{DF}_2$, which associates to a family of generalized K3-surfaces, a variation of Hodge structures.

In this setting, Theorem 1.1 should state that p induces isomorphisms of the Zariski tangent spaces of the two deformation functors. A generalization of this result to deformations of K3 surfaces of any order or to formal deformations would allow to prove that p is actually an isomorphism of functors.

To prove Theorem 1.1 we will need to show the compatibility between the actions of a Fourier–Mukai functor on Hochschild homology and singular cohomology which was conjecturally expected to hold true (see, for example, [8, 18]). Our result in this direction might be of independent interest and is the content of the following theorem, which will be proved in Section 2.2.

Theorem 1.2. *Let X_1 and X_2 be smooth complex projective varieties and let $\mathcal{E} \in \mathbf{D}^b(X_1 \times X_2)$. Then the following diagram*

$$\begin{array}{ccc} \mathrm{HH}_*(X_1) & \xrightarrow{(\Phi_{\mathcal{E}})_{\mathrm{HH}}} & \mathrm{HH}_*(X_2) \\ I_K^{X_1} \downarrow & & \downarrow I_K^{X_2} \\ H^*(X_1, \mathbb{C}) & \xrightarrow{(\Phi_{\mathcal{E}})_H} & H^*(X_2, \mathbb{C}) \end{array}$$

commutes.

The isomorphisms $I_K^{X_i}$ in the diagram above are modifications of the classical Hochschild–Kostant–Rosenberg isomorphisms by the square root of the Todd class of X_i (see [11, 18, 6]). A weaker version of Theorem 1.2 is proved in [6] for algebraic classes.

The paper ends with a discussion about the applications to the cases of Kummer and Enriques surfaces (see Section 4).

In this paper we will always work over the complex numbers. For X a smooth projective variety, $\mathbf{D}^b(X) := \mathbf{D}^b(\mathbf{Coh}(X))$.

2. HOCHSCHILD HOMOLOGY AND SINGULAR COHOMOLOGY

This section is mainly devoted to the proof of the commutativity of the actions of Fourier–Mukai equivalences on Hochschild homology and singular cohomology. To this end, we first show that such actions behave nicely with respect to the Künneth decomposition. The proof is then carried out using directly Ramadoss’ results [18, 25].

2.1. Hochschild homology and cohomology. Let X be a smooth projective variety and denote by ω_X its dualizing sheaf. Given the diagonal embedding $\Delta_X : X \hookrightarrow X \times X$, we define $S_X := \omega_X[\dim(X)]$, $S_X^{-1} := \omega_X^\vee[-\dim(X)]$ and $S_{\Delta_X}^{\pm 1} := (\Delta_X)_* S_X^{\pm 1}$. The i -th Hochschild homology and cohomology groups, $i \in \mathbb{Z}$, are now respectively (see, for example, [7])

$$\begin{aligned} \mathrm{HH}_i(X) &:= \mathrm{Hom}_{\mathbf{D}^b(X \times X)}(S_{\Delta_X}^{-1}[i], \mathcal{O}_{\Delta_X}) \cong \mathrm{Hom}_{\mathbf{D}^b(X)}(\mathcal{O}_X[i], \mathbf{L}\Delta_X^* \mathcal{O}_{\Delta_X}) \\ \mathrm{HH}^i(X) &:= \mathrm{Hom}_{\mathbf{D}^b(X \times X)}(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X}[i]) \cong \mathrm{Hom}_{\mathbf{D}^b(X)}(\mathbf{L}\Delta_X^* \mathcal{O}_{\Delta_X}, \mathcal{O}_X[i]). \end{aligned}$$

Hence one sets $\mathrm{HH}_*(X) := \bigoplus_i \mathrm{HH}_i(X)$ and $\mathrm{HH}^*(X) := \bigoplus_i \mathrm{HH}^i(X)$.

The Hochschild–Kostant–Rosenberg isomorphism $\mathbf{L}\Delta_X^* \mathcal{O}_{\Delta_X} \xrightarrow{\sim} \bigoplus_i \Omega_X^i[i]$ (see [6, 28]) yields the graded isomorphisms

$$I_{\mathrm{HKR}}^X : \mathrm{HH}_*(X) \rightarrow \mathrm{H}\Omega_*(X) := \bigoplus_i \mathrm{H}\Omega_i(X) \quad \text{and} \quad I_X^{\mathrm{HKR}} : \mathrm{HH}^*(X) \rightarrow \mathrm{HT}^*(X) := \bigoplus_i \mathrm{HT}^i(X),$$

where $\mathrm{H}\Omega_i(X) := \bigoplus_{q-p=i} H^p(X, \Omega_X^q)$ and $\mathrm{HT}^i(X) := \bigoplus_{p+q=i} H^p(X, \wedge^q \mathcal{T}_X)$. One then defines the graded isomorphisms

$$I_K^X = (\mathrm{td}(X)^{1/2} \wedge (-)) \circ I_{\mathrm{HKR}}^X \quad \text{and} \quad I_X^K = (\mathrm{td}(X)^{-1/2} \lrcorner (-)) \circ I_X^{\mathrm{HKR}}.$$

For X_1 and X_2 smooth and projective varieties and for $\mathcal{E} \in \mathrm{D}^b(X_1 \times X_2)$, let $\Phi_{\mathcal{E}} : \mathrm{D}^b(X_1) \rightarrow \mathrm{D}^b(X_2)$ be the *Fourier–Mukai functor* with kernel \mathcal{E} , i.e. the functor

$$\Phi_{\mathcal{E}} := \mathbf{R}(p_2)_*(\mathcal{E} \otimes^{\mathbf{L}} p_1^*(-)),$$

where $p_i : X_1 \times X_2 \rightarrow X_i$ is the i -th projection. The left and right adjoints of $\Phi_{\mathcal{E}}$ are again Fourier–Mukai functors with kernels $\mathcal{E}_L := \mathcal{E}^{\vee} \otimes p_2^* S_{X_2}$ and $\mathcal{E}_R := \mathcal{E}^{\vee} \otimes p_1^* S_{X_1}$ respectively (see [8, Prop. 5.9]). Notice that a Fourier–Mukai functor extends to a functor $\Phi_{\mathcal{E}} : \mathrm{D}(\mathbf{QCoh}(X_1)) \rightarrow \mathrm{D}(\mathbf{QCoh}(X_2))$.

From this, we get functorially a graded morphism $(\Phi_{\mathcal{E}})_{\mathrm{HH}} : \mathrm{HH}_*(X_1) \rightarrow \mathrm{HH}_*(X_2)$ such that, if $\mu \in \mathrm{HH}_i(X_1) = \mathrm{Hom}(S_{\Delta_{X_1}}^{-1}[i], \mathcal{O}_{\Delta_{X_1}})$, then $(\Phi_{\mathcal{E}})_{\mathrm{HH}}(\mu) \in \mathrm{HH}_i(X_2)$ is defined (see [7, Sect. 4.3]) by the composition

$$(2.1) \quad S_{\Delta_{X_2}}^{-1}[i] \xrightarrow{\gamma} \mathcal{E} \circ \mathcal{E}^{\vee}[i] \xrightarrow{\mathrm{id} \circ \eta \mathrm{id}} \mathcal{E} \circ S_{\Delta_{X_1}}^{-1}[i] \circ S_{\Delta_{X_1}} \circ \mathcal{E}^{\vee} \xrightarrow{\mathrm{id} \circ \mu \mathrm{id}} \mathcal{E} \circ S_{\Delta_{X_1}} \circ \mathcal{E}^{\vee} \xrightarrow{\circ} \mathcal{O}_{\Delta_{X_2}},$$

where $\mathcal{F} \circ \mathcal{G}$ is the kernel of the composition $\Phi_{\mathcal{F}} \circ \Phi_{\mathcal{G}}$ (see e.g. [8, Prop. 5.10]), $\eta : \mathcal{O}_{\Delta_{X_1}} \rightarrow S_{\Delta_{X_1}}^{-1} \circ S_{\Delta_{X_1}}$ is the isomorphism coming from the easy fact that $\Phi_{S_{\Delta_{X_1}}^{-1}} \circ \Phi_{S_{\Delta_{X_1}}} = \mathrm{id}$ and the morphisms γ and \circ are the natural ones (see [7, Appendix A]). If $\Phi_{\mathcal{E}}$ is an equivalence, then there exists also an action $(\Phi_{\mathcal{E}})^{\mathrm{HH}}$ on Hochschild cohomology induced by the functor $\Phi_{\mathcal{E} \boxtimes \mathcal{P}} : \mathrm{D}^b(X_1 \times X_1) \rightarrow \mathrm{D}^b(X_2 \times X_2)$, where $\mathcal{P} \cong \mathcal{E}_L \cong \mathcal{E}_R$ is the kernel of the inverse of $\Phi_{\mathcal{E}}$, which sends $\mathcal{O}_{\Delta_{X_1}}$ to $\mathcal{O}_{\Delta_{X_2}}$ (see, for example, [8, Remark 6.3]). The following easy result is probably well-known (see, for example, [27, Sect. 9.4] for a similar statement about Hochschild homology of rings).

Lemma 2.1. *Let X_1, X'_1, X_2 and X'_2 be smooth projective varieties.*

(i) *There exists a natural isomorphism*

$$\kappa_{X_1, X_2} : \mathrm{HH}_*(X_1) \otimes \mathrm{HH}_*(X_2) \rightarrow \mathrm{HH}_*(X_1 \times X_2), \quad (\alpha, \beta) \mapsto \alpha \boxtimes \beta := p_{X_1}^* \alpha \otimes p_{X_2}^* \beta$$

respecting the functoriality of Hochschild homology and compatible with the Hochschild–Kostant–Rosenberg isomorphism, i.e. the diagram

$$\begin{array}{ccc} \mathrm{HH}_*(X_1) \otimes \mathrm{HH}_*(X_2) & \xrightarrow{\kappa_{X_1, X_2}} & \mathrm{HH}_*(X_1 \times X_2) \\ I_{\mathrm{HKR}}^{X_1} \otimes I_{\mathrm{HKR}}^{X_2} \downarrow & & \downarrow I_{\mathrm{HKR}}^{X_1 \times X_2} \\ \mathrm{H}\Omega_*(X_1) \otimes \mathrm{H}\Omega_*(X_2) & \xrightarrow{\wedge} & \mathrm{H}\Omega_*(X_1 \times X_2) \end{array}$$

commutes.

(ii) *Given two Fourier–Mukai functors $\Phi_{\mathcal{E}} : \mathrm{D}^b(X_1) \rightarrow \mathrm{D}^b(X'_1)$ and $\Phi_{\mathcal{F}} : \mathrm{D}^b(X_2) \rightarrow \mathrm{D}^b(X'_2)$, we have*

$$(\Phi_{\mathcal{E} \boxtimes \mathcal{F}})_{\mathrm{HH}}(\mu \boxtimes \nu) = (\Phi_{\mathcal{E}})_{\mathrm{HH}}(\mu) \boxtimes (\Phi_{\mathcal{F}})_{\mathrm{HH}}(\nu),$$

for any $\mu \in \mathrm{HH}_(X_1)$ and $\nu \in \mathrm{HH}_*(X_2)$.*

Proof. Following [28], for any smooth projective variety X , the complex $\mathbf{L}\Delta_X^* \mathcal{O}_{\Delta_X}$ can be represented by the complex of \mathcal{O}_X -modules $\widehat{C}^{\bullet}(X)$ such that, if $q > 0$, then $\widehat{C}^q(X) = 0$, while, for $q \leq 0$ and for any affine $U = \mathrm{Spec}(R) \subseteq X$, the group of sections $\Gamma(U, \widehat{C}^q(X))$ is an adic completion of the usual module of Hochschild chains $C^q(R) = R^{\otimes -q+2} \otimes_{R \otimes R} R$. The differential is induced by the differential $d^n : R^{\otimes n+1} \rightarrow R^{\otimes n}$ such that

$$\begin{aligned} d^n(r_0 \otimes \dots \otimes r_n) &:= r_0 r_1 \otimes r_2 \otimes \dots \otimes r_n - r_0 \otimes r_1 r_2 \otimes \dots \otimes r_n + \dots \\ &\quad \dots + (-1)^{n-1} r_0 \otimes \dots \otimes r_{n-1} r_n + (-1)^n r_0 r_n \otimes \dots \otimes r_{n-1}. \end{aligned}$$

Moreover, the isomorphism I_{HKR}^X is given, locally in U , by the morphism $I^n(r_0 \otimes \dots \otimes r_n) = \frac{1}{n!} r_0 dr_1 \wedge \dots \wedge dr_n$ of R -modules.

For any pair of positive integers p, q , a (p, q) -shuffle is a permutation σ of $\{1, \dots, p+q\}$ such that $\sigma(1) < \dots < \sigma(p)$ and $\sigma(p+1) < \dots < \sigma(p+q)$. The morphism $\text{sh} : R_1^{\otimes p+1} \otimes_{\mathbb{C}} R_2^{\otimes q+1} \rightarrow (R_1 \otimes_{\mathbb{C}} R_2)^{\otimes p+q+1}$

$$\text{sh}((r_0 \otimes \dots \otimes r_p) \otimes_{\mathbb{C}} (r'_0 \otimes r_{p+1} \otimes \dots \otimes r_{p+q})) := \sum_{\sigma \in \text{Sh}(p,q)} (-1)^\sigma r_0 r'_0 \otimes r_{\sigma^{-1}(1)} \otimes \dots \otimes r_{\sigma^{-1}(p+q)},$$

where $\text{Sh}(p, q)$ is the set of all (p, q) -shuffles and X_i is locally $\text{Spec}(R_i)$, gives the local description of the isomorphisms $\kappa_{X,Y}$. The compatibility with the Hochschild–Kostant–Rosenberg isomorphism in (i) is now an easy check based on the previous local descriptions. (As pointed out by the referee, (i) could be probably proved directly by using the adjunction between the Hochschild–Kostant–Rosenberg isomorphism and the universal Atiyah class [6].)

To prove (ii), observe that $\Phi_{\mathcal{E} \boxtimes \mathcal{F}}(\mathcal{G}_1 \boxtimes \mathcal{G}_2) = \Phi_{\mathcal{E}}(\mathcal{G}_1) \boxtimes \Phi_{\mathcal{F}}(\mathcal{G}_2)$ and a similar decomposition holds true for the right and left adjoints. Since $\mathcal{O}_{\Delta_{X_1 \times X_2}} \cong \mathcal{O}_{\Delta_{X_1}} \boxtimes \mathcal{O}_{\Delta_{X_2}}$ and $S_{\Delta_{X_1 \times X_2}}^{-1} \cong S_{\Delta_{X_1}}^{-1} \boxtimes S_{\Delta_{X_2}}^{-1}$, the statement follows directly from the definition of the action of $\Phi_{\mathcal{E} \boxtimes \mathcal{F}}$ on Hochschild homology, since all the morphisms in (2.1) preserve the \boxtimes -product. \square

Remark 2.2. In the case of Fourier–Mukai equivalences, an analogous result can be proved for Hochschild cohomology by applying the same proof.

For later use, observe that any Fourier–Mukai functor $\Phi_{\mathcal{E}} : D^b(X_1) \rightarrow D^b(X_2)$ also induces an action on singular cohomology $(\Phi_{\mathcal{E}})_H : H^*(X_1, \mathbb{C}) \rightarrow H^*(X_2, \mathbb{C})$ by $(\Phi_{\mathcal{E}})_H(a) := (p_2)_*(v(\mathcal{E}) \cdot p_1^*(a))$, where $v(\mathcal{E}) := \text{ch}(\mathcal{E}) \cdot \sqrt{\text{td}(X_1 \times X_2)}$ is the *Mukai vector* of \mathcal{E} .

2.2. Proof of Theorem 1.2. For any smooth projective variety X , the Hochschild homology carries a non-degenerate pairing $\langle -, - \rangle_C : \text{HH}_i(X) \times \text{HH}_{-i}(X) \rightarrow \mathbb{C}$ such that, according to [7, Sect. 5], for any $\mu \in \text{HH}_i(X)$ and $\nu \in \text{HH}_{-i}(X)$,

$$\langle \mu, \nu \rangle_C := \text{tr}_{X \times X}(\tau(\mu) \circ \nu),$$

where $\tau : \mathbf{R}\text{Hom}_{X \times X}(S_{\Delta_X}^{-1}, \mathcal{O}_{\Delta_X}) \xrightarrow{\sim} \mathbf{R}\text{Hom}_{X \times X}(\mathcal{O}_{\Delta_X}, S_{\Delta_X})$ is the homomorphism obtained by tensoring on the right by $p_2^* S_X$, with $p_2 : X \times X \rightarrow X$ the second projection, and making the natural identifications.

If $\mathcal{E} \in D^b(X_1 \times X_2)$, then \mathcal{E} can be alternatively seen as a complex in $D^b(\text{pt} \times X_1 \times X_2)$ and hence there is a morphism $(\Phi_{\mathcal{E}})_{\text{HH}} : \text{HH}_0(\text{pt}) \rightarrow \text{HH}_0(X_1 \times X_2)$. Following [18, 7] one then defines the *Chern character of \mathcal{E}* as the element $\text{ch}(\mathcal{E}) := (\Phi_{\mathcal{E}})_{\text{HH}}(1) \in \text{HH}_0(X_1 \times X_2)$. The comparison with the standard Chern character in singular cohomology is the content of [6, Thm. 4.5] which yields $v(\mathcal{E}) = I_K^{X_1 \times X_2}(\text{ch}(\mathcal{E}))$.

With this in mind, one proves that the following equality holds true

$$(2.2) \quad \langle \text{ch}(\mathcal{E}^\vee), \mu \boxtimes \nu \rangle_C = \int_{X_2} I_K^{X_2}((\Phi_{\mathcal{E}})_{\text{HH}}(\mu)) \wedge I_K^{X_2}(\nu),$$

for any $\mu \in \text{HH}_*(X_1)$ and $\nu \in \text{HH}_*(X_2)$.

Indeed, first define the functor

$$\Psi_{\mathcal{E}} : D^b(X_1 \times X_2) \rightarrow D^b(\text{pt}), \quad \mathcal{F} \mapsto \mathbf{R}\Gamma(\mathcal{E} \otimes^{\mathbf{L}} \mathcal{F}).$$

An easy computation of the kernel of the composition of Fourier–Mukai functors shows that $\Psi_{\mathcal{E}}$ is isomorphic to the following composition

$$D^b(X_1 \times X_2) \xrightarrow{\Phi_{\mathcal{E}} \times \text{id}} D^b(X_2 \times X_2) \xrightarrow{\mathbf{L}\Delta_{X_2}^*} D^b(X_2) \xrightarrow{\mathbf{R}\Gamma} D^b(\text{pt}),$$

where $\Phi_{\mathcal{E}} \times \text{id} := \Phi_{\mathcal{E} \boxtimes \mathcal{O}_{\Delta_{X_2}}}$.

By definition $(\Psi_{\mathcal{E}})_{\text{HH}}(\mu \boxtimes \nu) = \langle 1, (\Psi_{\mathcal{E}})_{\text{HH}}(\mu \boxtimes \nu) \rangle_C$. By adjunction ([7, Thm. 8]) and the definition of Chern character, the latter is equal to $\langle \text{ch}(\mathcal{E}^{\vee}), \mu \boxtimes \nu \rangle_C$. Applying adjunction once more and Lemma 2.1(ii), we have

$$(\Psi_{\mathcal{E}})_{\text{HH}}(\mu \boxtimes \nu) = \langle \text{ch}(\mathcal{O}_{\Delta_{X_2}}^{\vee}), (\Phi_{\mathcal{E}} \times \text{id})_{\text{HH}}(\mu \boxtimes \nu) \rangle_C = \langle \text{ch}(\mathcal{O}_{\Delta_{X_2}}^{\vee}), (\Phi_{\mathcal{E}})_{\text{HH}}(\mu) \boxtimes \nu \rangle_C.$$

Following [25], denote by $K : \text{H}\Omega_*(X_i) \rightarrow \text{H}\Omega_*(X_i)$ the involution such that $K|_{H^q(X_i, \Omega_{X_i}^p)} = (-1)^q$. Using [25, Prop. 3] (or more precisely equation (8) in [25]) and Lemma 2.1(i), we get the following chain of equalities

$$\begin{aligned} \langle \text{ch}(\mathcal{O}_{\Delta_{X_2}}^{\vee}), (\Phi_{\mathcal{E}})_{\text{HH}}(\mu) \boxtimes \nu \rangle_C &= \int_{X_2 \times X_2} K(\text{ch}(\mathcal{O}_{\Delta_{X_2}}^{\vee})) \wedge I_{\text{HKR}}^{X_2 \times X_2}((\Phi_{\mathcal{E}})_{\text{HH}}(\mu) \boxtimes \nu) \wedge \text{td}(X_2 \times X_2) \\ &= \int_{X_2 \times X_2} \text{ch}(\mathcal{O}_{\Delta_{X_2}}) \wedge I_{\text{HKR}}^{X_2}((\Phi_{\mathcal{E}})_{\text{HH}}(\mu)) \wedge I_{\text{HKR}}^{X_2}(\nu) \wedge \text{td}(X_2 \times X_2) \\ &= \int_{X_2} I_K^{X_2}((\Phi_{\mathcal{E}})_{\text{HH}}(\mu)) \wedge I_K^{X_2}(\nu), \end{aligned}$$

for any $\mu \in \text{HH}_*(X_1)$ and $\nu \in \text{HH}_*(X_2)$. This proves (2.2).

Arguing in the same way, one shows that the following identities are true

$$\begin{aligned} \langle \text{ch}(\mathcal{E}^{\vee}), \mu \boxtimes \nu \rangle_C &= \int_{X_1 \times X_2} K(\text{ch}(\mathcal{E}^{\vee})) \wedge I_{\text{HKR}}^{X_1}(\mu) \wedge I_{\text{HKR}}^{X_2}(\nu) \wedge \text{td}(X_1 \times X_2) \\ &= \int_{X_1 \times X_2} v(\mathcal{E}) \wedge I_K^{X_1}(\mu) \wedge I_K^{X_2}(\nu) \\ &= \int_{X_2} (\Phi_{\mathcal{E}})_H(I_K^{X_1}(\mu)) \wedge I_K^{X_2}(\nu), \end{aligned}$$

for any $\mu \in \text{HH}_*(X_1)$ and $\nu \in \text{HH}_*(X_2)$. By (2.2), we have

$$\int_{X_2} I_K^{X_2}((\Phi_{\mathcal{E}})_{\text{HH}}(\mu)) \wedge I_K^{X_2}(\nu) = \int_{X_2} (\Phi_{\mathcal{E}})_H(I_K^{X_1}(\mu)) \wedge I_K^{X_2}(\nu),$$

for any $\mu \in \text{HH}_*(X_1)$ and $\nu \in \text{HH}_*(X_2)$.

From the fact that the pairing $\int_{X_2} (- \wedge -)$ is non-degenerate, $I_K^{X_2} \circ (\Phi_{\mathcal{E}})_{\text{HH}} = (\Phi_{\mathcal{E}})_H \circ I_K^{X_1}$. The theorem now follows from the natural identification $H^*(X, \mathbb{C}) \cong \text{H}\Omega_*(X)$ given by the Hodge decomposition.

Remark 2.3. As observed in [25], the singular cohomology groups $H^*(X_i, \mathbb{C})$ carry a non-degenerate pairing $\langle -, - \rangle_R : H^*(X_i, \mathbb{C}) \times H^*(X_i, \mathbb{C}) \rightarrow \mathbb{C}$ given by the formula (implicit in the previous proof)

$$\langle a, b \rangle_R := \int_{X_i} \frac{K(a)}{\sqrt{\text{ch}(\omega_{X_i})}} \wedge b,$$

for any $a, b \in H^*(X_i, \mathbb{C})$. If $\mathcal{E} \in \text{D}^b(X_1 \times X_2)$ is the kernel of a Fourier–Mukai equivalence, then the commutativity of the diagram in Theorem 1.2 and [25, Prop. 5] yields also the compatibility with the natural pairings, i.e.

$$\begin{aligned} \langle \alpha, \beta \rangle_C &= \langle I_K^{X_1}(\alpha), I_K^{X_1}(\beta) \rangle_R = \langle (\Phi_{\mathcal{E}})_{\text{HH}}(\alpha), (\Phi_{\mathcal{E}})_{\text{HH}}(\beta) \rangle_C \\ &= \langle I_K^{X_2}((\Phi_{\mathcal{E}})_{\text{HH}}(\alpha)), I_K^{X_2}((\Phi_{\mathcal{E}})_{\text{HH}}(\beta)) \rangle_R = \langle (\Phi_{\mathcal{E}})_H(I_K^{X_1}(\alpha)), (\Phi_{\mathcal{E}})_H(I_K^{X_1}(\beta)) \rangle_R \end{aligned}$$

for any $\alpha, \beta \in \text{HH}_*(X_1)$. The *generalized Mukai pairing* $\langle -, - \rangle_M : H^*(X_i, \mathbb{C}) \times H^*(X_i, \mathbb{C}) \rightarrow \mathbb{C}$, introduced in [6, Def. 3.2], is the non-degenerate pairing such that, for any $a, b \in H^*(X_i, \mathbb{C})$,

$$(2.3) \quad \langle a, b \rangle_M := \langle \tilde{\tau}(a), b \rangle_R$$

where $\tilde{\tau}|_{H^q(X, \Omega_X^p)} = (\sqrt{-1})^{-p-q}$. Such a pairing is also compatible with the action of $\Phi_{\mathcal{E}}$ on the singular cohomology.

3. INFINITESIMAL DEFORMATIONS

In this section we prove Theorem 1.1 which relates the existence of equivalences between some first order deformations of the derived categories of coherent sheaves and the existence of special Hodge isometries of the total cohomology. To this end, in Section 3.1 we briefly recall the description of the first order deformations of $\mathbf{Coh}(X)$ given in [26], for X a smooth projective variety. Notice that an equivalent theory can be obtained using the general results in [13, 15, 16]. Although the first approach is the preferred one in this paper, the latter will also be made use of at some specific points.

After this, in Section 3.2, we introduce a special weight-2 decomposition of the total cohomology groups (tensored by $\mathbb{Z}[\epsilon]/(\epsilon^2)$) preserved by the action of Fourier–Mukai equivalences.

3.1. The categorical setting. For X a smooth projective variety and $v \in \mathbb{H}^2(X)$, following [26], we consider the $\mathbb{C}[\epsilon]/(\epsilon^2)$ -linear abelian category $\mathbf{Coh}(X, v)$ which is the first order deformation of $\mathbf{Coh}(X)$ in the direction v . Since the precise definition of this category is not needed in the rest of this paper, we just recall the essentials of its construction.

Write $I_X^{\mathrm{HKR}}(v) = (\alpha, \beta, \gamma) \in \mathrm{HT}^2(X) = H^2(X, \mathcal{O}_X) \oplus H^1(X, \mathcal{T}_X) \oplus H^0(X, \wedge^2 \mathcal{T}_X)$. Then one defines a sheaf $\mathcal{O}_X^{(\beta, \gamma)}$ of $\mathbb{C}[\epsilon]/(\epsilon^2)$ -algebras on X depending only on β and γ . Representing $\alpha \in H^2(X, \mathcal{O}_X)$ as a Čech 2-cocycle $\{\alpha_{ijk}\}$ one has an element $\tilde{\alpha} := \{1 - \epsilon\alpha_{ijk}\}$ which is a Čech 2-cocycle with values in the invertible elements of the center of $\mathcal{O}_X^{(\beta, \gamma)}$. In analogy with the classical twisted setting, we get the abelian category $\mathbf{Coh}(\mathcal{O}_X^{(\beta, \gamma)}, \tilde{\alpha})$ of $\tilde{\alpha}$ -twisted coherent $\mathcal{O}_X^{(\beta, \gamma)}$ -modules. Now set $\mathbf{Coh}(X, v) := \mathbf{Coh}(\mathcal{O}_X^{(\beta, \gamma)}, \tilde{\alpha})$ and $\mathrm{D}^*(X, v) := \mathrm{D}^*(\mathbf{Coh}(X, v))$, where $*$ = b, \pm , \emptyset . Analogously, one defines the abelian category $\mathbf{QCoh}(X, v)$ of $\tilde{\alpha}$ -twisted quasi-coherent $\mathcal{O}_X^{(\beta, \gamma)}$ -modules, as the first order deformation of $\mathbf{QCoh}(X)$.

Remark 3.1. The construction of the abelian categories sketched above is a geometric incarnation of a more abstract theory developed in [16, 15]. The connection between the two approaches can be explained using [13], where the flat deformations of the abelian category of (quasi-)coherent sheaves on a variety are shown to be equivalent to flat deformations of the structure sheaf of X as a *twisted presheaf* (see [13, Thm. 1.4]). Such deformations are precisely the ones studied in [26].

The obstruction theory for lifting objects in these deformations has been developed in [14]. In particular, the obstruction class to deforming an object $\mathcal{E} \in \mathrm{D}^b(X)$ lives in $\mathrm{Ext}^2(\mathcal{E}, \mathcal{E})$, while all possible deformations form an affine space over $\mathrm{Ext}^1(\mathcal{E}, \mathcal{E})$.

Remark 3.2. According to [26, Sect. 4], the usual derived functors are well-defined in this context, provided the correct compatibilities. For example, consider the obvious morphism of algebras $\iota^\# : \mathcal{O}_X^{(\beta, \gamma)} \rightarrow \mathcal{O}_X$ which, for simplicity, will be denoted by ι . Then we have the functors $\iota_* : \mathrm{D}^b(X) \rightarrow \mathrm{D}^b(X, v)$, $\mathbf{L}\iota^* : \mathrm{D}^-(X, v) \rightarrow \mathrm{D}^-(X)$ and $\iota^! : \mathrm{D}^-(X, v) \rightarrow \mathrm{D}^-(X)$, where $\iota^!(-) := \bar{\iota}^* \mathbf{R}\mathcal{H}om_{\mathrm{D}^-(X, v)}(\iota_* \mathcal{O}_X, -)$ and $\bar{\iota}^*$ is induced by the natural exact functor between the categories of coherent $\iota_* \mathcal{O}_X$ -modules and $\mathbf{Coh}(X)$. The functors $\mathbf{L}\iota^*$ and $\iota^!$ are, respectively, left and right adjoints of ι_* .

Since there is no ambiguity, here as in the rest of the paper, we denote a functor $F : \mathrm{D}^b(X) \rightarrow \mathrm{D}^b(Y)$ and its (possible) extension to $\mathrm{D}^b(X, v) \rightarrow \mathrm{D}^b(Y, w)$ in the same way.

As observed in the introduction, by the definition of $\mathbf{Coh}(X, v)$, the notion of perfect complex makes sense also in this context. The category of perfect complexes is denoted by $\mathrm{D}_{\mathrm{perf}}(X, v)$. Take two smooth projective varieties X_1 and X_2 , $v_i \in \mathbb{H}^2(X_i)$ and $\tilde{\mathcal{E}} \in \mathrm{D}_{\mathrm{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$, where

$$-J(v_1) \boxplus v_2 := -p_1^* J(v_1) + p_2^* v_2$$

for $p_i : X_1 \times X_2 \rightarrow X_i$ the projection, and $J : \mathbb{H}^2(X_1) \rightarrow \mathbb{H}^2(X_1)$ is such that $(I_{X_1}^{\mathrm{HKR}} \circ J \circ (I_{X_1}^{\mathrm{HKR}})^{-1})(\alpha, \beta, \gamma) = (\alpha, -\beta, \gamma)$. (Notice that, when we write $-J(v_1) \boxplus v_2$, we are implicitly using

Remark 2.2.) Then the Fourier–Mukai functor

$$\Phi_{\tilde{\mathcal{E}}} : D^b(X_1, v_1) \rightarrow D^b(X_2, v_2)$$

is well-defined.

Proposition 3.3. *Let X_1 and X_2 be smooth projective varieties and $v_i \in \mathbb{H}^2(X_i)$. Take $\tilde{\mathcal{E}} \in D_{\text{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$ and set $\mathcal{E} := \mathbf{L}\iota^*\tilde{\mathcal{E}}$.*

(i) $\Phi_{\tilde{\mathcal{E}}} \circ \iota_* \cong \iota_* \circ \Phi_{\mathcal{E}} : D^b(X_1) \rightarrow D^b(X_2, v_2)$ and $\iota^! \circ \Phi_{\tilde{\mathcal{E}}} \cong \Phi_{\mathcal{E}} \circ \iota^! : D^b(X_1, v_1) \rightarrow D^-(X_2)$.

(ii) *If $\Phi_{\tilde{\mathcal{E}}} : D^b(X_1, v_1) \rightarrow D^b(X_2, v_2)$ is an equivalence, then $\Phi_{\mathcal{E}} : D^b(X_1) \rightarrow D^b(X_2)$ is an equivalence as well.*

Proof. The fact that $\Phi_{\tilde{\mathcal{E}}} \circ \iota_* \cong \iota_* \circ \Phi_{\mathcal{E}}$ was already remarked in [26, Thm. 4.7] and it is an easy application of the projection formula and flat base change (which in this context hold true as remarked in [26]). Indeed, for any $\mathcal{F} \in D^b(X_1)$,

$$\begin{aligned} \Phi_{\tilde{\mathcal{E}}}(\iota_*(\mathcal{F})) &= \mathbf{R}(p_2)_*(\tilde{\mathcal{E}} \otimes^{\mathbf{L}} p_1^*(\iota_*(\mathcal{F}))) \cong \mathbf{R}(p_2)_*(\tilde{\mathcal{E}} \otimes^{\mathbf{L}} \iota_*(p_1^*(\mathcal{F}))) \\ &\cong \mathbf{R}(p_2)_*\iota_*(\mathbf{L}\iota^*\tilde{\mathcal{E}} \otimes^{\mathbf{L}} p_1^*(\mathcal{F})) \cong \iota_*\mathbf{R}(p_2)_*(\mathcal{E} \otimes^{\mathbf{L}} p_1^*(\mathcal{F})) = \iota_*(\Phi_{\mathcal{E}}(\mathcal{F})). \end{aligned}$$

For the second one, we have that, for any $\mathcal{F} \in D^b(X_1, v_1)$,

$$\begin{aligned} \iota^!(\Phi_{\tilde{\mathcal{E}}}(\mathcal{F})) &= \iota^!\mathbf{R}(p_2)_*(\tilde{\mathcal{E}} \otimes^{\mathbf{L}} p_1^*(\mathcal{F})) \\ &\cong \mathbf{R}(p_2)_*(\iota^!(\tilde{\mathcal{E}} \otimes^{\mathbf{L}} p_1^*(\mathcal{F}))) \\ &\cong \mathbf{R}(p_2)_*(\mathcal{E} \otimes^{\mathbf{L}} \iota^!p_1^*(\mathcal{F})) \\ &\cong \mathbf{R}(p_2)_*(\mathcal{E} \otimes^{\mathbf{L}} p_1^*(\iota^!\mathcal{F})) \\ &= \Phi_{\mathcal{E}}(\iota^!(\mathcal{F})), \end{aligned}$$

where we used, as before, flat base change, the projection formula and the fact that $\tilde{\mathcal{E}}$ is a perfect complex.

To prove (ii), observe first that, as an easy consequence of [26, Cor. 3.3, Lemma 4.3], the category $\mathbf{QCoh}(X_i, v_i)$ has enough injectives. Moreover, $\iota^!\tilde{\mathcal{I}}$ is injective, for all injective objects $\tilde{\mathcal{I}} \in \mathbf{QCoh}(X_i, v_i)$, and the pull-backs of the injective objects in $\mathbf{QCoh}(X_i, v_i)$ via $\iota^!$ (co)generate $\mathbf{QCoh}(X_i)$ (actually all injective objects in $\mathbf{QCoh}(X_i)$ are of the form $\iota^!\tilde{\mathcal{I}}$, for some injective $\tilde{\mathcal{I}}$, by [16, Cor. 6.15]).

For $\mathcal{F} \in D^b(X_1)$ and an injective $\tilde{\mathcal{I}} \in D^b(X_1, v_1)$, we have the following chain of isomorphisms below

$$\begin{aligned} \text{Hom}(\mathcal{F}, \iota^!\tilde{\mathcal{I}}) &\cong \text{Hom}(\iota_*\mathcal{F}, \tilde{\mathcal{I}}) \\ &\cong \text{Hom}(\Phi_{\tilde{\mathcal{E}}}(\iota_*\mathcal{F}), \Phi_{\tilde{\mathcal{E}}}(\tilde{\mathcal{I}})) \\ &\cong \text{Hom}(\iota_*\Phi_{\mathcal{E}}(\mathcal{F}), \Phi_{\tilde{\mathcal{E}}}(\tilde{\mathcal{I}})) \\ &\cong \text{Hom}(\Phi_{\mathcal{E}}(\mathcal{F}), \iota^!(\Phi_{\tilde{\mathcal{E}}}(\tilde{\mathcal{I}}))) \\ &\cong \text{Hom}(\Phi_{\mathcal{E}}(\mathcal{F}), \Phi_{\mathcal{E}}(\iota^!\tilde{\mathcal{I}})), \end{aligned}$$

where the first and forth isomorphism are obtained by adjunction, the second one is simply the action of $\Phi_{\tilde{\mathcal{E}}}$, while the third and the last one are consequences of (i). It is easy to check that the composition of all these isomorphisms is the action of $\Phi_{\mathcal{E}}$. Since the objects $\iota^!\tilde{\mathcal{I}}$ (co)generate the category $\mathbf{QCoh}(X_1)$, the previous calculation shows that $\Phi_{\mathcal{E}}$ is fully-faithful.

Let $F : D^b(X_2, v_2) \rightarrow D^b(X_1, v_1)$ be a quasi-inverse of $\Phi_{\tilde{\mathcal{E}}}$. By (i) and adjunction, we have $F \circ \iota_* \cong \iota_* \circ \Phi_{\mathcal{E}_L}$. Thus, for any $\mathcal{F} \in D^b(X_2)$ and any injective $\tilde{\mathcal{I}} \in \mathbf{QCoh}(X_2, v_2)$,

$$\begin{aligned} \mathrm{Hom}(\mathcal{F}, \iota^! \tilde{\mathcal{I}}) &\cong \mathrm{Hom}(\iota_* \mathcal{F}, \tilde{\mathcal{I}}) \\ &\cong \mathrm{Hom}(F(\iota_* \mathcal{F}), F(\tilde{\mathcal{I}})) \\ &\cong \mathrm{Hom}(\iota_* \Phi_{\mathcal{E}_L}(\mathcal{F}), F(\tilde{\mathcal{I}})) \\ &\cong \mathrm{Hom}(\Phi_{\tilde{\mathcal{E}}}(\iota_* \Phi_{\mathcal{E}_L}(\mathcal{F})), \tilde{\mathcal{I}}) \\ &\cong \mathrm{Hom}(\iota_* \Phi_{\mathcal{E}}(\Phi_{\mathcal{E}_L}(\mathcal{F})), \tilde{\mathcal{I}}) \\ &\cong \mathrm{Hom}(\Phi_{\mathcal{E}}(\Phi_{\mathcal{E}_L}(\mathcal{F})), \iota^! \tilde{\mathcal{I}}). \end{aligned}$$

As before, the objects $\iota^! \tilde{\mathcal{I}}$ (co)generate the category $\mathbf{QCoh}(X_2)$. Therefore we have an isomorphism $\mathcal{F} \cong \Phi_{\mathcal{E}}(\Phi_{\mathcal{E}_L}(\mathcal{F}))$ which is what we wanted. An alternative proof can be obtained using Serre duality as explained in Appendix A. \square

The following result is the key ingredient in understanding the relation between first order deformations of varieties and deformations of kernels of Fourier–Mukai equivalences.

Theorem 3.4. (Toda) *Let X_1 and X_2 be smooth projective varieties and let $v_i \in \mathrm{HH}^2(X_i)$. Assume that there exists a Fourier–Mukai equivalence $\Phi_{\mathcal{E}} : D^b(X_1) \xrightarrow{\sim} D^b(X_2)$ with kernel $\mathcal{E} \in D^b(X_1 \times X_2)$. Then there exists an object $\tilde{\mathcal{E}} \in D_{\mathrm{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$ giving rise to a Fourier–Mukai equivalence $\Phi_{\tilde{\mathcal{E}}} : D^b(X_1, v_1) \xrightarrow{\sim} D^b(X_2, v_2)$ and such that $\mathcal{E} \cong \mathbf{L}\iota^* \tilde{\mathcal{E}}$ if and only if $(\Phi_{\mathcal{E}})^{\mathrm{HH}}(v_1) = v_2$.*

Proof. The existence of $\tilde{\mathcal{E}}$, given a Fourier–Mukai equivalence $\Phi_{\mathcal{E}}$ such that $(\Phi_{\mathcal{E}})^{\mathrm{HH}}(v_1) = v_2$, is precisely [26, Thm. 4.7]. For the proof of the other implication, we use an argument suggested to us by Y. Toda. Assume that $(\Phi_{\mathcal{E}})^{\mathrm{HH}}(v_1) = v_3$. Let $\mathcal{P} \in D^b(X_2 \times X_1)$ be the kernel of a quasi-inverse of $\Phi_{\mathcal{E}}$. By the first part of the theorem, there exists $\tilde{\mathcal{P}} \in D_{\mathrm{perf}}(X_2 \times X_1, -J(v_3) \boxplus v_1)$ giving rise to an equivalence $\Phi_{\tilde{\mathcal{P}}} : D^b(X_2, v_3) \rightarrow D^b(X_1, v_1)$. The composition $G := \Phi_{\tilde{\mathcal{E}}} \circ \Phi_{\tilde{\mathcal{P}}}$ induces an equivalence $\mathbf{Coh}(X_2, v_3) \rightarrow \mathbf{Coh}(X_2, v_2)$ (as deformations of $\mathbf{Coh}(X_2)$). Indeed, by Proposition 3.3(i), $G(\iota_* \mathbf{Coh}(X_2)) \subseteq \iota_* \mathbf{Coh}(X_2)$. Since the abelian categories $\mathbf{Coh}(X_2, v_2)$ and $\mathbf{Coh}(X_2, v_3)$ are generated by $\iota_* \mathbf{Coh}(X_2)$ by extensions, G yields the desired equivalence. But now, by [15, Thm. 3.1], all the deformations of $\mathbf{Coh}(X_2)$ are parametrized by $\mathrm{HH}^2(X_2)$. Thus $v_2 = v_3$. \square

3.2. Infinitesimal Mukai lattices and the proof of Theorem 1.1. Let X be a K3 surface and let $v \in \mathrm{HH}^2(X)$. Let $w := I_K^X(\sigma_X) + \epsilon I_K^X(v \circ \sigma_X) \in \tilde{H}(X, \mathbb{Z}) \otimes \mathbb{C}[\epsilon]/(\epsilon^2)$, where σ_X is a generator for $\mathrm{HH}_2(X)$ as a \mathbb{C} -vector space. Here $\tilde{H}(X, \mathbb{Z})$ is the *Mukai lattice* of X , i.e. the \mathbb{Z} -module $H^*(X, \mathbb{Z})$ endowed with the (generalized) Mukai pairing and the weight-2 Hodge structure

$$\tilde{H}^{2,0}(X) := H^{2,0}(X) \quad \tilde{H}^{0,2}(X) := \overline{\tilde{H}^{2,0}(X)} \quad \tilde{H}^{1,1}(X) := (\tilde{H}^{2,0}(X) \oplus \tilde{H}^{0,2}(X))^{\perp}.$$

Definition 3.5. The free $\mathbb{Z}[\epsilon]/(\epsilon^2)$ -module of finite rank $\tilde{H}(X, \mathbb{Z}) \otimes \mathbb{Z}[\epsilon]/(\epsilon^2)$ endowed with the $\mathbb{Z}[\epsilon]/(\epsilon^2)$ -linear extension of the generalized Mukai pairing $\langle -, - \rangle_M$ and such that $\tilde{H}(X, \mathbb{Z}) \otimes \mathbb{C}[\epsilon]/(\epsilon^2)$ has the weight-2 decomposition

$$\begin{aligned} \tilde{H}^{2,0}(X, v) &:= \mathbb{C}[\epsilon]/(\epsilon^2) \cdot w \\ \tilde{H}^{0,2}(X, v) &:= \overline{\tilde{H}^{2,0}(X, v)} \\ \tilde{H}^{1,1}(X, v) &:= (\tilde{H}^{2,0}(X, v) \oplus \tilde{H}^{0,2}(X, v))^{\perp}, \end{aligned}$$

is the *infinitesimal Mukai lattice* of X with respect to v , which is denoted by $\tilde{H}(X, v, \mathbb{Z})$.

Remark 3.6. (i) It is easy to see that w behaves like a honest period of a K3 surface. More precisely,

$$\langle w, w \rangle_M = 0 \quad \langle w, \bar{w} \rangle_M > 0,$$

where we do not distinguish between the Mukai pairing (2.3) and its $\mathbb{Z}[\epsilon]/(\epsilon^2)$ -linear extension. Hence the weight-2 decomposition in the previous definition can be thought of as the analogue of the weight-2 Hodge decomposition on $\tilde{H}(X, \mathbb{Z})$ appearing in the classical Derived Torelli Theorem (see, for example, [8] for the classical case and [10] for the twisted setting).

(ii) Going back to the motivation described in the introduction, suppose the morphism $p : \mathrm{DF}_1 \rightarrow \mathrm{DF}_2$ of functors is given. Theorem 1.1 would then state the equivariance of the differential of p with respect to the action of Fourier–Mukai equivalences on the first order versions of the functors DF_1 and DF_2 .

Given this analogy, for two K3 surfaces X_1 and X_2 , and $v_i \in \mathrm{HH}^2(X_i)$, a *Hodge isometry* of the infinitesimal Mukai lattices is a $\mathbb{Z}[\epsilon]/(\epsilon^2)$ -linear isomorphism $g : \tilde{H}(X_1, v_1, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(X_2, v_2, \mathbb{Z})$ preserving the Mukai pairing and the weight-2 decomposition in the previous definition. In the rest of this paper, we will be more interested in the infinitesimal isometries $g = g_0 \otimes \mathbb{Z}[\epsilon]/(\epsilon^2)$, where g_0 is (automatically) an Hodge isometry of the standard Mukai lattices $\tilde{H}(X_1, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(X_2, \mathbb{Z})$. These special infinitesimal Hodge isometries will be called *effective*.

The lattice $\tilde{H}(X_i, \mathbb{Z})$ has some interesting substructures. Indeed, let σ_i be a generator of $H^{2,0}(X_i)$ and ω_i a Kähler class. Then

$$(3.1) \quad P(X_i, \sigma_i, \omega_i) := \langle \mathrm{Re}(\sigma_i), \mathrm{Im}(\sigma_i), 1 - \omega_i^2/2, \omega_i \rangle,$$

is a positive four-space in $\tilde{H}(X_i, \mathbb{R})$ (here $\mathrm{Re}(\sigma_i)$ and $\mathrm{Im}(\sigma_i)$ are the real and imaginary part of σ_i). It comes, by the choice of basis, with a natural orientation. An effective Hodge isometry $g = g_0 \otimes \mathbb{Z}[\epsilon]/(\epsilon^2) : \tilde{H}(X_1, v_1, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(X_2, v_2, \mathbb{Z})$ is *orientation preserving* if g_0 preserves the orientation of $P(X, \sigma_i, \omega_i)$, for $i = 1, 2$. For $X = X_1 = X_2$ and $v = v_1 = v_2$, the group of orientation preserving effective Hodge isometries is denoted by $O_+(\tilde{H}(X, v, \mathbb{Z}))$.

Proof of Theorem 1.1. We first prove that (i) implies (ii). Let $\Phi_{\tilde{\mathcal{E}}} : \mathrm{D}^b(X_1, v_1) \xrightarrow{\sim} \mathrm{D}^b(X_2, v_2)$ be an equivalence with kernel $\tilde{\mathcal{E}} \in \mathrm{D}_{\mathrm{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$. By Proposition 3.3(ii), $\mathcal{E} := \mathbf{L}i^* \tilde{\mathcal{E}} \in \mathrm{D}^b(X_1 \times X_2)$ is the kernel of a Fourier–Mukai equivalence $\Phi_{\mathcal{E}} : \mathrm{D}^b(X_1) \xrightarrow{\sim} \mathrm{D}^b(X_2)$. Corollary 7.9 in [9] implies that the Hodge isometry $g_0 := (\Phi_{\mathcal{E}})_H : \tilde{H}(X_1, \mathbb{Z}) \rightarrow \tilde{H}(X_2, \mathbb{Z})$ is orientation preserving. Moreover, by Theorem 3.4, since $\tilde{\mathcal{E}}$ is a first order deformation of \mathcal{E} , $(\Phi_{\mathcal{E}})^{\mathrm{HH}}(v_1) = v_2$.

Consider the diagram

$$(3.2) \quad \begin{array}{ccc} \mathrm{HH}^*(X_1) & \xrightarrow{(\Phi_{\mathcal{E}})^{\mathrm{HH}}} & \mathrm{HH}^*(X_2) \\ (-) \circ \sigma_{X_1} \downarrow & & \downarrow (-) \circ (\Phi_{\mathcal{E}})_{\mathrm{HH}}(\sigma_{X_1}) \\ \mathrm{HH}_*(X_1) & \xrightarrow{(\Phi_{\mathcal{E}})_{\mathrm{HH}}} & \mathrm{HH}_*(X_2) \\ I_K^{X_1} \downarrow & & \downarrow I_K^{X_2} \\ \tilde{H}(X_1, \mathbb{C}) & \xrightarrow{(\Phi_{\mathcal{E}})_H} & \tilde{H}(X_2, \mathbb{C}), \end{array}$$

where, as before, σ_{X_1} is a generator of $\mathrm{HH}_2(X_1)$ as a \mathbb{C} -vector space and $(-) \circ (-) : \mathrm{HH}^*(X_i) \times \mathrm{HH}_*(X_i) \rightarrow \mathrm{HH}_*(X_i)$ denotes the action of $\mathrm{HH}^*(X_i)$ on $\mathrm{HH}_*(X_i)$ (see, for example, [7]). The upper square is commutative by, for example, [8, Remark 6.3], while the commutativity of the bottom one is Theorem 1.2. Thus

$$g_0(I_K^{X_1}(v_1 \circ \sigma_{X_1})) = I_K^{X_2}(v_2 \circ ((I_K^{X_1})^{-1} \circ g_0 \circ I_K^{X_1})(\sigma_{X_1})).$$

In particular, $g := g_0 \otimes \mathbb{Z}[\epsilon]/(\epsilon^2) : \tilde{H}(X_1, v_1, \mathbb{Z}) \rightarrow \tilde{H}(X_2, v_2, \mathbb{Z})$ is an effective orientation preserving Hodge isometry of the infinitesimal Mukai lattices.

The fact that (ii) implies (i) is shown as follows. Let $g = g_0 \otimes \mathbb{Z}[\epsilon]/(\epsilon^2)$ be as in (ii), with $g_0 : \tilde{H}(X_1, \mathbb{Z}) \rightarrow \tilde{H}(X_2, \mathbb{Z})$ an orientation preserving Hodge isometry. By the classical Derived Torelli Theorem [19, 22], there exists a Fourier–Mukai equivalence $\Phi_{\mathcal{E}} : D^b(X_1) \xrightarrow{\sim} D^b(X_2)$ with kernel $\mathcal{E} \in D^b(X_1 \times X_2)$ and such that $g_0 = (\Phi_{\mathcal{E}})_H$.

Under our assumptions, the commutativity of diagram (3.2) gives $(\Phi_{\mathcal{E}})^{\mathrm{HH}}(v_1) = v_2$. Therefore, by Theorem 3.4, there exists a first order deformation $\tilde{\mathcal{E}} \in D_{\mathrm{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$ of \mathcal{E} such that the Fourier–Mukai functor $\Phi_{\tilde{\mathcal{E}}} : D^b(X_1, v_1) \xrightarrow{\sim} D^b(X_2, v_2)$ is an equivalence. \square

Theorem 1.1 is precisely the classical result by Mukai and Orlov [19, 22] if v_1 and v_2 in the statement are trivial. In particular, under this assumption, condition (ii) can be relaxed avoiding the orientation preserving requirement. This is no longer true when v_1 and v_2 are non-trivial, as explained in the example below.

Example 3.7. Let X be a K3 surface with $\mathrm{Pic}(X) = \mathbb{Z}H$ and $H^2 > 2$. Take $v \in \mathrm{HH}^2(X)$ such that $w := I_K^X(v \circ \sigma_X) = (\sqrt{2}, \sqrt{3}H, \sqrt{5}) \in \tilde{H}^{1,1}(X)$, where σ_X is a generator of $\mathrm{HH}_2(X)$. The Hodge isometry $j := \mathrm{id}_{(H^0 \oplus H^4)(X, \mathbb{Z})} \oplus (-\mathrm{id}_{H^2(X, \mathbb{Z})})$ does not preserve the orientation and maps w to $w' := (\sqrt{2}, -\sqrt{3}H, \sqrt{5})$. Let $v' \in \mathrm{HH}^2(X)$ be such that $w' = I_K^X(v' \circ \sigma_X)$. Then $D^b(X, v)$ is not Fourier–Mukai equivalent to $D^b(X, v')$.

Indeed, suppose for a contradiction that they are Fourier–Mukai equivalent. By Theorem 1.1, there is an orientation preserving Hodge isometry g of $\tilde{H}(X, \mathbb{Z})$ such that $g(v) = \alpha v'$, for some root of unity α . But

$$g(\sqrt{2}, \sqrt{3}H, \sqrt{5}) = (a_1\sqrt{2} + a_2\sqrt{3} + a_3\sqrt{5}, (b_1\sqrt{2} + b_2\sqrt{3} + b_3\sqrt{5})H, c_1\sqrt{2} + c_2\sqrt{3} + c_3\sqrt{5}),$$

with $a_i, b_i, c_i \in \mathbb{Z}$, $i = 1, 2, 3$. An easy computation shows that $g(v) = \alpha v'$ only if g restricted to $H^0(X, \mathbb{Z}) \oplus \mathrm{Pic}(X) \oplus H^4(X, \mathbb{Z})$ is such that $g(x, yH, z) = \pm(x, -yH, z)$. By [21, Lemma 4.1], $g|_{T(X)} = \pm \mathrm{id}$ and hence by [20, Thm. 1.6.1, Cor. 1.5.2], $g = \pm j$, which is a contradiction.

Notice that in such a case, no object in $D^b(X)$ deforms to an object in $D^b(X, v)$, or to one in $D^b(X, v')$.

Remark 3.8. Given a K3 surface X , the number of isomorphism classes of K3 surfaces with derived category equivalent to $D^b(X)$ is finite (see [4]). The same result holds true in the broader case of twisted K3 surfaces (see [10, Cor. 4.6]).

As first order deformations of X are parametrized by an affine space over $\mathrm{HH}^2(X)$ this cannot be true in the deformed setting. Hence one could weaken the notion of isomorphism between deformed K3 surfaces following [10], requiring that (X_1, v_1) and (X_2, v_2) (where X_i is a K3 surface and $v_i \in \mathrm{HH}^2(X_i)$) are *equivalent deformations* if there exists an isomorphism $f : X_1 \rightarrow X_2$ such that $f^*v_2 = v_1$. Unfortunately, the number of Fourier–Mukai partners of a deformed K3 surface (X, v) remains infinite even for this equivalence relation.

Indeed, let X be a K3 surface containing infinitely many smooth rational curves $\{C_i\}_{i \in \mathbb{N}}$ and take $v \in \mathrm{HH}^2(X)$ such that $w := I_K^X(v \circ \sigma_X) = (1, H, 1) \in \tilde{H}^{1,1}(X)$, for $H \in \mathrm{Pic}(X)$ an ample line bundle and σ_X a generator of $\mathrm{HH}_2(X)$. If s_i is the Hodge isometry of the total cohomology group of X which acts as the reflection in the class of the rational curve C_i , then for any $r \in \mathbb{N}$ $w_r := (s_r \circ \dots \circ s_1)(w)$ yields pairwise non-equivalent deformations (X, v_r) , with $v_r \in \mathrm{HH}^2(X)$, such that $D^b(X, v) \cong D^b(X, v_r)$ (by Theorem 1.1). The same example shows that the number of Fourier–Mukai partners is infinite, even if we declare the deformations (X_1, v_1) and (X_2, v_2) (with X_i and $v_i \in \mathrm{HH}^2(X_i)$ as before) to be isomorphic if there exists an equivalence $(L \otimes (-)) \circ f^* : \mathbf{Coh}(X_2, v_2) \xrightarrow{\sim} \mathbf{Coh}(X_1, v_1)$, for some isomorphism $f : X_1 \xrightarrow{\sim} X_2$ and some line bundle $L \in \mathrm{Pic}(X_1)$.

3.3. Fourier–Mukai functors and the group of autoequivalences. In this section we generalize a few properties of Fourier–Mukai functors to the case of deformed categories with the aim of specializing Theorem 1.1 to autoequivalences.

Take X_1 , X_2 and X_3 smooth projective varieties and $v_i \in \mathbb{H}^2(X_i)$, where $i = 1, 2, 3$ and $I_{X_i}^{\text{HKR}}(v_i) = (\alpha_i, \beta_i, \gamma_i)$. Let $\tilde{\mathcal{E}} \in \text{D}_{\text{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$ and $\tilde{\mathcal{F}} \in \text{D}_{\text{perf}}(X_2 \times X_3, -J(v_2) \boxplus v_3)$. Define

$$\tilde{\mathcal{F}} \circ \tilde{\mathcal{E}} := \mathbf{R}(p_{13})_*(p_{12}^* \tilde{\mathcal{E}} \otimes^{\mathbf{L}} p_{23}^* \tilde{\mathcal{F}}) \in \text{D}^b(X_1 \times X_3, -J(v_1) \boxplus v_3),$$

where p_{ij} are the natural projections from $X_1 \times X_2 \times X_3$. To unravel the definition, observe that, for $\{i, j\} \in \{\{1, 2\}, \{2, 3\}\}$, one has $p_{ij}^* : \text{D}^b(X_i \times X_j, -J(v_i) \boxplus v_j) \rightarrow \text{D}^b(X_1 \times X_2 \times X_3, w_{ij})$, where

$$\begin{aligned} I_{X_1 \times X_2 \times X_3}^{\text{HKR}}(w_{12}) &= (p_{12}^*(-\alpha_1 \boxplus \alpha_2), \beta_1 \boxplus \beta_2 \boxplus \beta_3, -\gamma_1 \boxplus \gamma_2 \boxplus 0) \\ I_{X_1 \times X_2 \times X_3}^{\text{HKR}}(w_{23}) &= (p_{23}^*(-\alpha_2 \boxplus \alpha_3), \beta_1 \boxplus \beta_2 \boxplus \beta_3, 0 \boxplus -\gamma_2 \boxplus \gamma_3). \end{aligned}$$

Moreover, we can tensor $p_{12}^* \tilde{\mathcal{E}}$ and $p_{23}^* \tilde{\mathcal{F}}$, seen respectively as objects in the derived categories of $q_2^{-1} \mathcal{O}_{X_2}^{(\beta_2, \gamma_2)}$ -modules and $q_2^{-1} \mathcal{O}_{X_2}^{(\beta_2, -\gamma_2)}$ -modules, where $q_2 : X_1 \times X_2 \times X_3 \rightarrow X_2$ is the projection. Such a tensor product takes naturally values in the derived category of $p_{13}^{-1} \mathcal{O}_{X_1 \times X_3}^{(\beta_1 \boxplus \beta_3, -\gamma_1 \boxplus \gamma_3)}$ -modules. Hence, by [26], we can apply the functor $\mathbf{R}(p_{13})_*$. In this argument we did not take care of the twist because it behaves nicely with respect to the various operations.

Lemma 3.9. *Under the above assumptions, $\tilde{\mathcal{G}} := \tilde{\mathcal{F}} \circ \tilde{\mathcal{E}} \in \text{D}_{\text{perf}}(X_1 \times X_3, -J(v_1) \boxplus v_3)$.*

Proof. Let $\mathcal{E} := \mathbf{L}^* \tilde{\mathcal{E}}$ and $\mathcal{F} := \mathbf{L}^* \tilde{\mathcal{F}}$. To prove that $\tilde{\mathcal{G}}$ is perfect it is sufficient to show that $\mathbf{L}^* \tilde{\mathcal{G}}$ is bounded. This is an easy consequence of the following isomorphisms

$$\begin{aligned} \mathbf{L}^* \tilde{\mathcal{G}} &= \mathbf{L}^* \mathbf{R}(p_{13})_*(p_{12}^* \tilde{\mathcal{E}} \otimes^{\mathbf{L}} p_{23}^* \tilde{\mathcal{F}}) \\ &\cong \mathbf{R}(p_{13})_* \mathbf{L}^*(p_{12}^* \tilde{\mathcal{E}} \otimes^{\mathbf{L}} p_{23}^* \tilde{\mathcal{F}}) \\ &\cong \mathbf{R}(p_{13})_* \mathbf{L}(p_{12}^* \mathcal{E} \otimes^{\mathbf{L}} p_{23}^* \mathcal{F}) \\ &= \mathcal{F} \circ \mathcal{E}, \end{aligned}$$

where the natural isomorphism $\mathbf{L}^* \mathbf{R}(p_{13})_* \cong \mathbf{R}(p_{13})_* \mathbf{L}^*$ was already observed in the proof of Lemma 6.5 in [26]. Such an isomorphism will be further clarified in Appendix A (see Lemma A.5). \square

Due to this result, the composition of the Fourier–Mukai functors $\Phi_{\tilde{\mathcal{F}}}$ and $\Phi_{\tilde{\mathcal{E}}}$ is again a Fourier–Mukai functor with kernel $\tilde{\mathcal{F}} \circ \tilde{\mathcal{E}}$. In the case of non-deformed derived categories, the inverse of any Fourier–Mukai equivalence is again of Fourier–Mukai type. This fact needs to be proved in the first order deformation case.

Proposition 3.10. *If X_1 and X_2 are smooth projective varieties, $v_i \in \mathbb{H}^2(X_i)$ and*

$$\Phi_{\tilde{\mathcal{E}}} : \text{D}^b(X_1, v_1) \xrightarrow{\sim} \text{D}^b(X_2, v_2)$$

is a Fourier–Mukai equivalence with $\tilde{\mathcal{E}} \in \text{D}_{\text{perf}}(X_1 \times X_2, -J(v_1) \boxplus v_2)$, then the inverse is a Fourier–Mukai functor.

Proof. This result can be easily proved using Serre duality as in Appendix A (see Corollary A.8). Nevertheless we can also argue as follows. Observe first that, as an easy application of the projection formula, the identity $\text{D}^b(X_i, v_i) \rightarrow \text{D}^b(X_i, v_i)$ is a Fourier–Mukai functor whose kernel is $(\Delta_{X_i})_* \mathcal{O}_{X_i}^{(\beta_i, \gamma_i)}$, where $I_{X_i}^{\text{HKR}}(v_i) = (\alpha_i, \beta_i, \gamma_i)$. Notice that $\mathbf{L}^*(\Delta_{X_i})_* \mathcal{O}_{X_i}^{(\beta_i, \gamma_i)} \cong \mathcal{O}_{\Delta_{X_i}}$.

Let $\mathcal{P} \in \mathrm{D}^b(X_2 \times X_1)$ be the kernel of the inverse of the equivalence $\Phi_{\mathcal{E}}$ (see Proposition 3.3(ii)), where $\mathcal{E} := \mathbf{L}\iota^*\tilde{\mathcal{E}}$. By Theorem 3.4, there exists at least a $\tilde{\mathcal{P}} \in \mathrm{D}_{\mathrm{perf}}(X_2 \times X_1, -J(v_2) \boxplus v_1)$ such that $\mathcal{P} \cong \mathbf{L}\iota^*\tilde{\mathcal{P}}$. By [14], all such kernels are parametrized by $\mathrm{Ext}^1(\mathcal{P}, \mathcal{P})$.

The functor $\tilde{\mathcal{E}} \circ (-) : \mathrm{D}_{\mathrm{perf}}(X_2 \times X_1, -J(v_2) \boxplus v_1) \rightarrow \mathrm{D}_{\mathrm{perf}}(X_2 \times X_2, -J(v_2) \boxplus v_2)$ induces an isomorphism $\mathrm{Ext}^1(\mathcal{P}, \mathcal{P}) \xrightarrow{\sim} \mathrm{Ext}^1(\mathcal{O}_{\Delta_{X_2}}, \mathcal{O}_{\Delta_{X_2}})$ and hence a one-to-one correspondence between deformations of \mathcal{P} and $\mathcal{O}_{\Delta_{X_2}}$. Therefore, by the previous computation, there exists a $\tilde{\mathcal{P}}$ such that $\tilde{\mathcal{E}} \circ \tilde{\mathcal{P}} \cong (\Delta_{X_2})_* \mathcal{O}_{X_2}^{(\beta_2, \gamma_2)}$. \square

As a consequence, for a smooth projective variety X and $v \in \mathrm{HH}^2(X)$, the set $\mathrm{Aut}^{\mathrm{FM}}(\mathrm{D}^b(X, v))$ of all autoequivalences of Fourier–Mukai type of $\mathrm{D}^b(X, v)$ is actually a group.

As remarked in the introduction, when X is a K3 surface, Theorem 1.1 can be read in terms of the existence of a surjective group homomorphism

$$\Pi_{(X,v)} : \mathrm{Aut}^{\mathrm{FM}}(\mathrm{D}^b(X, v)) \longrightarrow \mathrm{O}_+(\tilde{H}(X, v, \mathbb{Z})),$$

where $\mathrm{O}_+(\tilde{H}(X, v, \mathbb{Z}))$ denotes the group of orientation preserving effective Hodge isometries. If $v = 0$, the kernel of $\Pi_{(X,0)}$ consists of all autoequivalences acting trivially on cohomology and, according to Conjecture 1.2 in [2], should be described as the fundamental group of a period domain naturally associated to a connected component of the manifold parametrizing stability conditions on $\mathrm{D}^b(X)$.

The relation with the case $v \neq 0$ is clarified by the following easy result.

Lemma 3.11. *If X is a K3 surface and $v \in \mathrm{HH}^2(X)$, then $\ker(\Pi_{(X,v)}) \cong \ker(\Pi_{(X,0)})$.*

Proof. Any autoequivalence $\Phi_{\tilde{\mathcal{E}}}$ is in $\ker(\Pi_{(X,v)})$ if and only if $\Pi_{(X,v)}(\Phi_{\tilde{\mathcal{E}}}) = \mathrm{id} \otimes \mathbb{Z}[\epsilon]/(\epsilon^2)$. In this case, by Proposition 3.3(ii) $\mathcal{E} := \mathbf{L}\iota^*\tilde{\mathcal{E}}$ is the kernel of a Fourier–Mukai equivalence which, by the proof of Theorem 1.1, acts trivially on cohomology. Hence, there exists a morphism $\kappa : \ker(\Pi_{(X,v)}) \rightarrow \ker(\Pi_{(X,0)})$ sending $\tilde{\mathcal{E}}$ to \mathcal{E} , which is surjective by Theorem 3.4. By [14], given $\mathcal{E} \in \mathrm{D}^b(X \times X)$, all the $\tilde{\mathcal{E}} \in \mathrm{D}_{\mathrm{perf}}(X \times X, -J(v) \boxplus v)$ such that $\mathcal{E} = \mathbf{L}\iota^*\tilde{\mathcal{E}}$ form an affine space over $\mathrm{Ext}^1(\mathcal{E}, \mathcal{E})$ which, in the case of K3 surfaces, is trivial. Thus κ is an isomorphism. \square

4. FURTHER EXAMPLES

Let X be a smooth projective variety with an action of a finite group G . We denote by $\mathbf{Coh}_G(X)$ the abelian category of G -equivariant coherent sheaves on X , i.e. the category whose objects are pairs $(\mathcal{E}, \{\lambda_g\}_{g \in G})$, where $\mathcal{E} \in \mathbf{Coh}(X)$ and, for any $g_1, g_2 \in G$, $\lambda_{g_i} : \mathcal{E} \xrightarrow{\sim} g_i^*\mathcal{E}$ is an isomorphism such that $\lambda_{g_1 g_2} = g_2^*(\lambda_{g_1}) \circ \lambda_{g_2}$. The set of these isomorphisms is a G -linearization of \mathcal{E} (very often a G -linearization will be simply denoted by λ). The morphisms in $\mathbf{Coh}_G(X)$ are just the morphisms of coherent sheaves compatible with the G -linearizations. We put $\mathrm{D}_G^b(X) := \mathrm{D}^b(\mathbf{Coh}_G(X))$. Since G is finite, $\mathrm{D}_G^b(X)$ can equivalently be described in terms of G -equivariant objects in $\mathrm{D}^b(X)$ (see, for example, [24, Sect. 1.1]). For later use, we recall the definition of the functor $\mathrm{Inf}_G : \mathrm{D}^b(X) \rightarrow \mathrm{D}_G^b(X)$

$$\mathrm{Inf}_G(\mathcal{E}) := \left(\bigoplus_{g \in G} g^*\mathcal{E}, \lambda_{\mathrm{nat}} \right),$$

where λ_{nat} is the natural G -linearization.

4.1. Kummer surfaces. Let now A be an abelian surface and denote by $\mathrm{Km}(A)$ the corresponding Kummer surface, i.e. the minimal resolution of the quotient of A by the natural involution $\varpi : A \rightarrow A$, with $\varpi(a) = -a$. Denote by $G \cong \mathbb{Z}/2\mathbb{Z}$ the group generated by ϖ . The main result in [3]

shows that there exists a Fourier–Mukai equivalence $\Psi_A : D_G^b(A) \xrightarrow{\sim} D^b(\mathrm{Km}(A))$. The composition $\Pi_A := \Psi_A \circ \mathrm{Inf}_G : D^b(A) \rightarrow D^b(\mathrm{Km}(A))$ is of Fourier–Mukai type and induces a morphism

$$(\Pi_A)_{\mathrm{HH}} : \mathrm{HH}_*(A) \rightarrow \mathrm{HH}_*(\mathrm{Km}(A)).$$

Thus, given $v \in \mathrm{HH}^2(A)$, we get $\pi(v, \sigma_A) := (\Pi_A)_{\mathrm{HH}}(v \circ \sigma_A) \circ (\Pi_A)_{\mathrm{HH}}(\sigma_A)^{-1} \in \mathrm{HH}^2(\mathrm{Km}(A))$, where σ_A is a generator of $\mathrm{HH}_2(A)$.

Just as the classical Derived Torelli Theorem does, the infinitesimal version in Theorem 1.1 holds true for abelian surfaces as well. From this we deduce a relation between the deformations of Fourier–Mukai equivalences in the case of abelian surfaces and the ones for the corresponding Kummer surfaces.

Proposition 4.1. *Let A_1 and A_2 be abelian surfaces and let $v_i \in \mathrm{HH}^2(A_i)$, with $i = 1, 2$, be such that there exists a Fourier–Mukai equivalence*

$$\Phi_{\tilde{\mathcal{E}}} : D^b(A_1, v_1) \xrightarrow{\sim} D^b(A_2, v_2).$$

Then $D^b(\mathrm{Km}(A_1), \pi(v_1, \sigma_{A_1}))$ and $D^b(\mathrm{Km}(A_2), \pi(v_2, (\Phi_{\mathcal{E}})_{\mathrm{HH}}(\sigma_{A_1})))$ are Fourier–Mukai equivalent, where $\mathcal{E} := \mathbf{L}t^*\tilde{\mathcal{E}} \in D^b(A_1 \times A_2)$.

Proof. The same argument as in [17, Prop. 3.4] shows that there exists $\mathcal{G} \in D^b(A_1 \times A_2)$ giving rise to an equivalence $\Phi_{\mathcal{G}} : D^b(A_1) \xrightarrow{\sim} D^b(A_2)$ such that $(\varpi \times \varpi)^*\mathcal{G} \cong \mathcal{G}$ and $(\Phi_{\mathcal{G}})_H = (\Phi_{\mathcal{E}})_H$. Hence, by Theorem 1.2, the fact that $(\Phi_{\mathcal{G}})_{\mathrm{HH}} = (\Phi_{\mathcal{E}})_{\mathrm{HH}}$ and the commutativity of the diagram

$$\begin{array}{ccc} \mathrm{HH}^*(A_1) & \xrightarrow{(\Phi_{\mathcal{G}})^{\mathrm{HH}}} & \mathrm{HH}^*(A_2) \\ (-) \circ \sigma_{A_1} \downarrow & & \downarrow (-) \circ (\Phi_{\mathcal{G}})_{\mathrm{HH}}(\sigma_{A_1}) \\ \mathrm{HH}_*(A_1) & \xrightarrow{(\Phi_{\mathcal{G}})_{\mathrm{HH}}} & \mathrm{HH}_*(A_2), \end{array}$$

we have $(\Phi_{\mathcal{G}})^{\mathrm{HH}}(v_1) = v_2$.

By the discussion in [24, Sect. 3.2] and Theorem 1.2, there exists an object $\mathcal{F} \in D^b(\mathrm{Km}(A_1) \times \mathrm{Km}(A_2))$ inducing an equivalence $\Phi_{\mathcal{F}} : D^b(\mathrm{Km}(A_1)) \xrightarrow{\sim} D^b(\mathrm{Km}(A_2))$ and making commutative the following diagram

$$\begin{array}{ccccc} \mathrm{HH}_*(A_1) & \xrightarrow{(\Phi_{\mathcal{G}})_{\mathrm{HH}}} & & \mathrm{HH}_*(A_2) & \\ \downarrow (\Pi_{A_1})_{\mathrm{HH}} & \searrow I_K^{A_1} & \tilde{H}(A_1, \mathbb{C}) \xrightarrow{(\Phi_{\mathcal{G}})_H} \tilde{H}(A_2, \mathbb{C}) & \swarrow I_K^{A_2} & \downarrow (\Pi_{A_2})_{\mathrm{HH}} \\ & & \downarrow & & \\ & & \tilde{H}(\mathrm{Km}(A_1), \mathbb{C}) \xrightarrow{(\Phi_{\mathcal{F}})_H} \tilde{H}(\mathrm{Km}(A_2), \mathbb{C}) & & \\ & \swarrow I_K^{\mathrm{Km}(A_1)} & & \swarrow I_K^{\mathrm{Km}(A_2)} & \\ \mathrm{HH}_*(\mathrm{Km}(A_1)) & \xrightarrow{(\Phi_{\mathcal{F}})_{\mathrm{HH}}} & & \mathrm{HH}_*(\mathrm{Km}(A_2)) & \end{array}$$

The commutativity of (3.2), for $X_i = \mathrm{Km}(A_i)$, and Theorem 3.4 yield the following chain of equalities

$$\begin{aligned}
 (\Phi_{\mathcal{F}})^{\mathrm{HH}}(\pi(v_1, \sigma_{A_1})) &= (\Phi_{\mathcal{F}})^{\mathrm{HH}}((\Pi_{A_1})_{\mathrm{HH}}(v_1 \circ \sigma_{A_1}) \circ (\Pi_{A_1})_{\mathrm{HH}}(\sigma_{A_1})^{-1}) \\
 &= (\Phi_{\mathcal{F}})_{\mathrm{HH}}((\Pi_{A_1})_{\mathrm{HH}}(v_1 \circ \sigma_{A_1})) \circ ((\Phi_{\mathcal{F}})_{\mathrm{HH}}((\Pi_{A_1})_{\mathrm{HH}}(\sigma_{A_1})))^{-1} \\
 &= (\Pi_{A_2})_{\mathrm{HH}}((\Phi_{\mathcal{G}})_{\mathrm{HH}}(v_1 \circ \sigma_{A_1})) \circ ((\Pi_{A_2})_{\mathrm{HH}}((\Phi_{\mathcal{G}})_{\mathrm{HH}}(\sigma_{A_1})))^{-1} \\
 &= (\Pi_{A_2})_{\mathrm{HH}}(v_2 \circ (\Phi_{\mathcal{G}})_{\mathrm{HH}}(\sigma_{A_1})) \circ ((\Pi_{A_2})_{\mathrm{HH}}((\Phi_{\mathcal{G}})_{\mathrm{HH}}(\sigma_{A_1})))^{-1} \\
 &= (\Pi_{A_2})_{\mathrm{HH}}(v_2 \circ (\Phi_{\mathcal{E}})_{\mathrm{HH}}(\sigma_{A_1})) \circ ((\Pi_{A_2})_{\mathrm{HH}}((\Phi_{\mathcal{E}})_{\mathrm{HH}}(\sigma_{A_1})))^{-1} \\
 &= \pi(v_2, (\Phi_{\mathcal{E}})_{\mathrm{HH}}(\sigma_{A_1})).
 \end{aligned}$$

Theorem 3.4 concludes the proof. \square

In general, even when we consider deformations of Kummer surfaces induced by those of the corresponding abelian surfaces, the converse of the previous result it is not expected to hold true.

4.2. Enriques surfaces. Let Y be an Enriques surface, i.e. a minimal smooth projective surface with 2-torsion canonical bundle ω_Y and $H^1(Y, \mathcal{O}_Y) = 0$. The universal cover $\pi : X \rightarrow Y$ is a K3 surface and it carries a fixed-point-free involution $\varpi : X \rightarrow X$ such that $Y = X/G$, where $G = \langle \varpi \rangle$. In this special setting, $\mathbf{Coh}(Y)$ is naturally isomorphic to the abelian category $\mathbf{Coh}_G(X)$ which yields an equivalence $\mathrm{D}^b(Y) \cong \mathrm{D}_G^b(X)$, which will be tacitly meant for what follows.

Notice that, by functoriality, since π is an étale morphism, we have an induced morphism $\pi^* : \mathrm{HH}^*(Y) \rightarrow \mathrm{HH}^*(X)$ which is compatible with the Hochschild–Kostant–Rosenberg isomorphism.

Proposition 4.2. *Let Y_1 and Y_2 be Enriques surfaces, $\pi_i : X_i \rightarrow Y_i$ be their universal covers, and $v_i \in \mathrm{HH}^2(Y_i)$, for $i = 1, 2$. Then the following are equivalent:*

(i) *There exists a Fourier–Mukai equivalence*

$$\Phi_{\tilde{\mathcal{E}}} : \mathrm{D}^b(Y_1, v_1) \xrightarrow{\sim} \mathrm{D}^b(Y_2, v_2)$$

with $\tilde{\mathcal{E}} \in \mathrm{D}_{\mathrm{perf}}(Y_1 \times Y_2, -J(v_1) \boxplus v_2)$.

(ii) *There exists an orientation preserving effective Hodge isometry*

$$g : \tilde{H}(X_1, \pi_1^*(v_1), \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(X_2, \pi_2^*(v_2), \mathbb{Z})$$

which is G -equivariant, i.e. $\varpi^ \circ g = g \circ \varpi^*$.*

(iii) *There exists a Fourier–Mukai equivalence*

$$\Phi_{\tilde{\mathcal{F}}} : \mathrm{D}^b(X_1, \pi_1^*(v_1)) \xrightarrow{\sim} \mathrm{D}^b(X_2, \pi_2^*(v_2))$$

with $\tilde{\mathcal{F}} \in \mathrm{D}_{\mathrm{perf}}(X_1 \times X_2, -J(\pi_1^(v_1)) \boxplus \pi_2^*(v_2))$ such that $(\varpi \times \varpi)^* \mathcal{F} \cong \mathcal{F}$, where $\mathcal{F} := \mathbf{L}t^* \tilde{\mathcal{F}}$.*

Proof. The equivalence of (ii) and (iii) is simply a rewriting of Theorem 1.1 in the equivariant context using [17, Prop. 3.4] (or better its version for equivalences). Now, due to [24, Sect. 3.3], the existence of the kernel $\mathcal{F} \in \mathrm{D}^b(X_1 \times X_2)$ of a Fourier–Mukai equivalence $\Phi_{\mathcal{F}}$ such that $(\varpi \times \varpi)^* \mathcal{F} \cong \mathcal{F}$ is equivalent to the existence of an object $\mathcal{E} \in \mathrm{D}^b(Y_1 \times Y_2)$ which gives rise to an equivalence $\Phi_{\mathcal{E}} : \mathrm{D}^b(Y_1) \xrightarrow{\sim} \mathrm{D}^b(Y_2)$. Hence, the equivalence of (i) and (iii) is a consequence of the commutativity of the following diagram

$$\begin{array}{ccc}
 \mathrm{HH}^*(Y_1) & \xrightarrow{(\Phi_{\mathcal{E}})^{\mathrm{HH}}} & \mathrm{HH}^*(Y_2) \\
 \pi_1^* \downarrow & & \downarrow \pi_2^* \\
 \mathrm{HH}^*(X_1) & \xrightarrow{(\Phi_{\mathcal{F}})^{\mathrm{HH}}} & \mathrm{HH}^*(X_2).
 \end{array}$$

This commutativity can be checked using the isomorphism of functors $\pi_2^* \circ \Phi_{\mathcal{E}} \cong \Phi_{\mathcal{F}} \circ \pi_1^* : \mathrm{D}^b(Y_1) \rightarrow \mathrm{D}^b(X_2)$, which in turn can be deduced from [8, Sect. 7.3] and [24]. \square

Acknowledgements. The final version of this paper was written during the authors' stay at the Hausdorff Center for Mathematics (Bonn) whose hospitality is gratefully acknowledged. It is a pleasure to thank Daniel Huybrechts for useful advice, Wendy Lowen and Yukinobu Toda for kindly answering our questions and the referee for helping us clarifying the motivations of the main result of this paper. We are grateful to Sukhendu Mehrotra for taking part in the discussion in which the main idea of this paper came to light and for writing the appendix. The second named author was partially supported by the Università degli Studi di Milano (FIRST 2004).

REFERENCES

- [1] O. Ben-Bassat, J. Block, T. Pantev, *Non-commutative tori and Fourier–Mukai duality*, *Compositio Math.* **143** (2007), 423–475.
- [2] T. Bridgeland, *Stability conditions on K3 surfaces*, *Duke Math. J.* **141** (2008), 241–291.
- [3] T. Bridgeland, A. King, M. Reid, *The McKay correspondence as an equivalence of derived categories*, *J. Am. Math. Soc.* **14** (2001), 535–554.
- [4] T. Bridgeland, A. Maciocia, *Complex surfaces with equivalent derived categories*, *Math. Z.* **236** (2001), 677–697.
- [5] D. Burns, M. Rapoport, *On the Torelli problem for Kählerian K-3 surfaces*, *Ann. Sci. cole Norm. Sup.* **8** (1975), 235–273.
- [6] A. Căldăraru, *The Mukai pairing II: The Hochschild–Kostant–Rosenberg isomorphism*, *Adv. Math.* **194** (2005), 34–66.
- [7] A. Căldăraru, S. Willerton, *The Mukai pairing, I: a categorical approach*, arXiv:0707.2052.
- [8] D. Huybrechts, *Fourier–Mukai transforms in algebraic geometry*, Oxford Mathematical Monographs (2006).
- [9] D. Huybrechts, E. Macrì, P. Stellari, *Derived equivalences of K3 surfaces and orientation*, to appear in: *Duke Math. J.* arXiv:0710.1645.
- [10] D. Huybrechts, P. Stellari, *Equivalences of twisted K3 surfaces*, *Math. Ann.* **332** (2005), 901–936.
- [11] M. Kontsevich, *Deformation quantization of Poisson manifolds*, *Lett. Math. Phys.* **66** (2003), 157–216.
- [12] E. Looijenga, P.Ch. Peters, *Torelli theorems for Kähler K3 surfaces*, *Compositio Math.* **42** (1980/81), 145–186.
- [13] W. Lowen, *Algebroid prestacks and deformations of ringed spaces*, *Trans. Amer. Math. Soc.* **360** (2008), 1631–1660.
- [14] W. Lowen, *Obstruction theory for objects in abelian and derived categories*, *Comm. Algebra* **33** (2005), 3195–3223.
- [15] W. Lowen, M. Van den Bergh, *Hochschild cohomology of abelian categories and ringed spaces*, *Adv. Math.* **198** (2005), 172–221.
- [16] W. Lowen, M. Van den Bergh, *Deformation theory of abelian categories*, *Trans. Amer. Math. Soc.* **358** (2006), 5441–5483.
- [17] E. Macrì, S. Mehrotra, P. Stellari, *Inducing stability conditions*, to appear in: *J. Algebraic Geom.* arXiv:0705.3752.
- [18] N. Markarian, *Poincaré–Birkhoff–Witt isomorphism, Hochschild homology and Riemann–Roch theorem*. Preprint MPI 2001-52.
- [19] S. Mukai, *On the moduli space of bundles on K3 surfaces, I*, In: *Vector Bundles on Algebraic Varieties*, Oxford University Press, Bombay and London (1987), 341–413.
- [20] V.V. Nikulin, *Integral symmetric bilinear forms and some of their applications*, *Math. USSR Izvestija* **14** (1980), 103–167.
- [21] K. Oguiso, *K3 surfaces via almost-primes*, *Math. Res. Lett.* **9** (2002), 47–63.
- [22] D. Orlov, *Equivalences of derived categories and K3 surfaces*, *J. Math. Sci.* **84** (1997), 1361–1381.
- [23] I. Pijateckij–Šapiro, I. Šafarevič, *A Torelli theorem for algebraic surfaces of type K3*, *Math. USSR Izvestija* **5** (1971), 547–588.
- [24] D. Ploog, *Equivariant autoequivalences for finite group actions*, *Adv. Math.* **216** (2007), 62–74.
- [25] A. Ramadoss, *The relative Riemann–Roch theorem from Hochschild homology*, *New York J. Math.* **14** (2008), 643–717.
- [26] Y. Toda, *Deformations and Fourier–Mukai transforms*, *J. Diff. Geom.* **81** (2009), 197–224.
- [27] C. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics 38, Cambridge University Press (1994).
- [28] A. Yekutieli, *The continuous Hochschild cochain complex of a scheme*, *Canadian J. Math.* **54** (2002), 1319–1337.

APPENDIX A. DUALITY FOR INFINITESIMAL DEFORMATIONS

by SUKHENDU MEHROTRA

Let X, Y be smooth and projective varieties, with $(0, \beta, \gamma) \in \text{HT}^2(X)$, $(0, \beta', \gamma') \in \text{HT}^2(Y)$, and suppose $f : (X, \mathcal{O}_X^{(\beta, \gamma)}) \rightarrow (Y, \mathcal{O}_Y^{(\beta', \gamma')})$ is a morphism of locally ringed spaces over $R_1 := \mathbb{C}[\epsilon]/(\epsilon^2)$ (the notation here is that of Section 3.1). Then, for $\alpha' \in H^2(Y, \mathcal{O}_Y)$, there is defined a push-forward functor between twisted derived categories $\mathbf{R}f_* : \text{D}^b(\mathbf{Coh}(\mathcal{O}_X^{(\beta, \gamma)}, \widetilde{f^*\alpha'})) \rightarrow \text{D}^b(\mathbf{Coh}(\mathcal{O}_Y^{(\beta', \gamma')}, \widetilde{\alpha'}))$ ([7]). The purpose of this appendix is to prove the existence of a right adjoint $f^! : \text{D}^b(\mathbf{Coh}(\mathcal{O}_Y^{(\beta', \gamma')}, \widetilde{\alpha'})) \rightarrow \text{D}^b(\mathbf{Coh}(\mathcal{O}_X^{(\beta, \gamma)}, \widetilde{f^*\alpha'}))$ to $\mathbf{R}f_*$ under suitable hypotheses on f . More precisely, what is proven is the existence of a “dualizing complex.”

The setting here will be slightly more general than that of the main article. To begin with, let X denote a separated, finite type scheme over \mathbb{C} . By an *infinitesimal deformation* of X , we will mean a locally ringed space $\widetilde{X} = (X, \mathcal{A}_X)$, where \mathcal{A}_X is a sheaf of local flat R_1 -algebras, with a fixed isomorphism $\mathcal{A}_X \otimes_{R_1} \mathbb{C} \cong \mathcal{O}_X$, such that

$$(*) \quad 0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{A}_X \rightarrow \mathcal{O}_X \rightarrow 0$$

is a central extension. In addition, we require that the following condition be satisfied:

- (**) for any local section $f \in \mathcal{O}_X(U)$ over an affine open set U , there is a lift $\hat{f} \in \mathcal{A}_X(U)$ such that the multiplicative set $\{\hat{f}^n : n \geq 0\}$ satisfies the left and right Ore localization conditions (see [4], 2.1.6).

Naturally, infinitesimal deformations then form a category in the obvious way.

If X is affine, and M is a $\mathcal{A}_X(X)$ module, we define the *sheaf associated to M* as the sheaf M^\sim which on principal opens X_f has sections $M_{\hat{f}}$. A quick word about this construction: If $X_g \subset X_f$, we have that $g^n = f\alpha$, for some $n > 0$ and $\alpha \in \mathcal{O}_X(X)$. Thus, by condition (*), \hat{g}^n and $\hat{f}\hat{\alpha}$ differ by a central square-zero element, so that if the first is invertible, so is the other. The universal property of localization ([4]) then yields a restriction map $M_{\hat{f}} \rightarrow M_{\hat{g}}$; the uniqueness of this map shows that as defined, M^\sim is indeed a sheaf. The alert reader will have noticed that we tacitly assumed a *choice* of a lift \hat{f} for any given $f \in \mathcal{O}_X(X)$ in making this definition. However, the same reasoning as above shows that any other choice of lifts yields a canonically isomorphic sheaf. Finally, let us mention that an important property of this construction is that the assignment $M \mapsto M^\sim$ is an exact functor (see propositions 2.1.16 (ii) and 2.1.17 (i) of [4]).

The categories $\mathbf{Coh}(\widetilde{X}) := \mathbf{Coh}(\mathcal{A}_X)$ and $\mathbf{QCoh}(\widetilde{X}) := \mathbf{QCoh}(\mathcal{A}_X)$ of coherent and quasi-coherent sheaves of *right* modules are defined in the usual way, in terms of free presentations on open sets. Using the exactness of the associated-sheaf construction, it can easily be verified that the analogue of Lemma 3.1 in [7] holds: to wit, for any $\mathcal{F} \in \mathbf{QCoh}(\widetilde{X})$ and any affine open $U \subset X$, $\mathcal{F}|_U = \mathcal{F}(U)^\sim$, the sheaf associated to the module $\mathcal{F}(U)$, and that the functor $M \mapsto M^\sim$ gives an equivalence $\mathcal{A}_{\widetilde{X}}(U)\text{-Mod} \rightarrow \mathbf{QCoh}(\widetilde{U})$.

It is standard that the category $\mathcal{A}_{\widetilde{X}}\text{-Mod}$ of $\mathcal{A}_{\widetilde{X}}$ -modules has enough injective and flat objects ([7]). So, for any morphism $f : \widetilde{X} \rightarrow \widetilde{Y}$ of infinitesimal deformations, the functors $f_* : \mathcal{A}_{\widetilde{X}}\text{-Mod} \rightarrow \mathcal{A}_{\widetilde{Y}}\text{-Mod}$ and $f^* : \mathcal{A}_{\widetilde{Y}}\text{-Mod} \rightarrow \mathcal{A}_{\widetilde{X}}\text{-Mod}$ between categories of right modules extend to derived functors $\mathbf{R}f_* : \text{D}^+(\mathcal{A}_{\widetilde{X}}\text{-Mod}) \rightarrow \text{D}^+(\mathcal{A}_{\widetilde{Y}}\text{-Mod})$ and $\mathbf{L}f^* : \text{D}^-(\mathcal{A}_{\widetilde{Y}}\text{-Mod}) \rightarrow \text{D}^-(\mathcal{A}_{\widetilde{X}}\text{-Mod})$. In fact, for our purposes, we shall employ the general result Theorem 4.5 of [6] from which it follows that $\mathbf{R}f_*$ extends to a functor $\mathbf{R}f_* : \text{D}(\mathcal{A}_{\widetilde{X}}\text{-Mod}) \rightarrow \text{D}(\mathcal{A}_{\widetilde{Y}}\text{-Mod})$ between *unbounded* derived categories. Subtleties like the equivalence between $\text{D}\mathbf{QCoh}(\mathcal{A}_{\widetilde{X}}\text{-Mod})$ and $\text{D}(\mathbf{QCoh}(\widetilde{X}))$ have been dealt with in [1] and are not a cause for concern; it is then immediately seen that $\mathbf{R}f_*$ and $\mathbf{L}f^*$ are defined between categories of quasi-coherent sheaves.

The next couple of results establish the existence of a right adjoint to the pushforward functor for unbounded derived categories of quasi-coherent sheaves. This is a direct application of Neeman's general approach to duality via topological methods.

Lemma A.1. *Let $f : \tilde{X} \rightarrow \tilde{Y}$ be a morphism of infinitesimal deformations, with X a separated scheme and $\bar{f} := f \otimes_{R_1} \mathbb{C}$ a separated morphism of schemes. Then, the functor $\mathbf{R}f_* : \mathbf{D}(\mathbf{QCoh}(\tilde{X})) \rightarrow \mathbf{D}(\mathbf{QCoh}(\tilde{Y}))$ respects coproducts, that is, for any small set Λ , the natural map $\coprod_{\lambda \in \Lambda} \mathbf{R}f_*(\mathcal{F}_\lambda) \rightarrow \mathbf{R}f_*(\coprod_{\lambda \in \Lambda} \mathcal{F}_\lambda)$ is an isomorphism.*

Proof. The proof of Lemma 1.4 in [5] carries over word-for-word. \square

Theorem A.2. (Neeman) *Under the hypotheses of the previous lemma, $\mathbf{R}f_*$ admits a right adjoint $f^! : \mathbf{D}(\mathbf{QCoh}(\tilde{Y})) \rightarrow \mathbf{D}(\mathbf{QCoh}(\tilde{X}))$.*

Proof. The proofs of Lemma 2.6 in [5] and Proposition 6.1 in [1] can be adapted to our setting to show that if $\mathcal{F} \in \mathbf{D}(\mathbf{QCoh}(\tilde{X}))$ satisfies $\mathrm{Hom}_{\tilde{X}}(\mathcal{L}, \mathcal{F}) = 0$ for every perfect object \mathcal{L} , it must in fact be the zero object; in other words, the category $\mathbf{D}(\mathbf{QCoh}(\tilde{X}))$ is compactly generated. This, and the previous lemma, furnish the hypotheses of Theorem 4.1 in [5], from which the result follows. \square

In the following paragraphs, we prove the claimed coherence and boundedness properties of the right adjoint $f^!$. From now on, all morphisms between infinitesimal deformations will be assumed to be of *finite type*.

Denote by $\iota_X : X \rightarrow \tilde{X}$ and $\iota_Y : Y \rightarrow \tilde{Y}$ the canonical immersions; we observe that the functors $(\iota_X)_*$ and $(\iota_Y)_*$ are exact.

Lemma A.3. *Let $f : \tilde{X} \rightarrow \tilde{Y}$ be a flat morphism of infinitesimal deformations, with $\bar{f} : X \rightarrow Y$ a smooth morphism of schemes. Then given $\mathcal{F} \in \mathbf{D}(\mathbf{QCoh}(Y))$, there is an isomorphism $(\iota_X)_* \bar{f}^! \mathcal{F} \cong f^! (\iota_Y)_* \mathcal{F}$ in $\mathbf{D}(\mathbf{QCoh}(\tilde{X}))$.*

Assuming the previous lemma for the moment, we have the following result.

Proposition A.4. *Let $f : \tilde{X} \rightarrow \tilde{Y}$ be a flat morphism of infinitesimal deformations, with \bar{f} smooth. Then, $f^! : \mathbf{D}(\mathbf{QCoh}(\tilde{Y})) \rightarrow \mathbf{D}(\mathbf{QCoh}(\tilde{X}))$ restricts to a functor $f^! : \mathbf{D}_{\mathrm{perf}}(\tilde{Y}) \rightarrow \mathbf{D}^b(\mathbf{Coh}(\tilde{X}))$, the category $\mathbf{D}_{\mathrm{perf}}(\tilde{Y})$ being that of perfect complexes on \tilde{Y} .*

Proof. Given $\mathcal{F} \in \mathbf{D}_{\mathrm{perf}}(\tilde{Y})$, one has the exact triangle:

$$(\iota_Y)_* \mathbf{L}\iota_Y^* \mathcal{F} \xrightarrow{\epsilon} \mathcal{F} \rightarrow (\iota_Y)_* \mathbf{L}\iota_Y^* \mathcal{F}.$$

Applying the functor $f^!$ and rearranging, using Lemma A.3, we get the exact triangle:

$$(\iota_X)_* \bar{f}^! (\mathbf{L}\iota_Y^* \mathcal{F}) \rightarrow f^! \mathcal{F} \rightarrow (\iota_X)_* \bar{f}^! (\mathbf{L}\iota_Y^* \mathcal{F}).$$

As $\bar{f}^!$ exists between derived categories of *coherent* sheaves and $\mathbf{L}\iota_Y^* \mathcal{F} \in \mathbf{D}^b(\mathbf{Coh}(\tilde{X}))$, the outer two terms are in $\mathbf{D}^b(\mathbf{Coh}(\tilde{X}))$. Consequently, the middle term is also. \square

Proof of Lemma A.3. Pick any object $\mathcal{G} \in \mathbf{D}^-(\mathbf{QCoh}(\tilde{X}))$ and $\mathcal{F} \in \mathbf{D}(\mathbf{QCoh}(Y))$. It follows by adjunction that

$$\mathbf{R}\mathrm{Hom}_{\tilde{X}}(\mathcal{G}, (\iota_X)_* \bar{f}^! \mathcal{F}) \cong \mathbf{R}\mathrm{Hom}_{\tilde{Y}}(\mathbf{R}\bar{f}_* \mathbf{L}\iota_X^* \mathcal{G}, \mathcal{F}).$$

Assume the base-change property which will be proved in Lemma A.5: $\mathbf{L}\iota_Y^* \mathbf{R}f_* \cong \mathbf{R}\bar{f}_* \mathbf{L}\iota_X^*$, that is, the standard natural transformation of functors is an isomorphism. Then, continuing from the previous line, we find that

$$\begin{aligned} \mathbf{R}\mathrm{Hom}_{\tilde{X}}(\mathcal{G}, (\iota_X)_* \bar{f}^! \mathcal{F}) &\cong \mathbf{R}\mathrm{Hom}_{\tilde{Y}}(\mathbf{L}\iota_Y^* \mathbf{R}f_* \mathcal{G}, \mathcal{F}) \\ &\cong \mathbf{R}\mathrm{Hom}_{\tilde{X}}(\mathcal{G}, f^! (\iota_Y)_* \mathcal{F}), \end{aligned}$$

all isomorphisms being canonical and functorial in the first argument; call their composition *can*. To simplify typography, set $\mathcal{A} = (\iota_X)_* \bar{f}^! \mathcal{F}$ and $\mathcal{B} := f^! (\iota_Y)_* \mathcal{F}$. Express \mathcal{A} as the homotopy colimit (see Remark 2.2, [1]) :

$$\coprod_{i>0} (\mathcal{A})^{\leq i} \xrightarrow{id-s} \coprod_{i>0} (\mathcal{A})^{\leq i} \longrightarrow \mathcal{A}$$

where the bounded above complex $(\mathcal{A})^{\leq i}$ is the i -th truncation:

$$\mathcal{H}^n \left((\mathcal{A})^{\leq i} \right) = \begin{cases} \mathcal{H}^n(\mathcal{A}), & \text{if } n \leq i \\ 0, & \text{if } n > i, \end{cases}$$

and s is the direct sum of the morphisms $(\mathcal{A})^{\leq i} \rightarrow (\mathcal{A})^{\leq i+1}$. This yields the following diagram with exact triangles for rows:

$$\begin{array}{ccccc} \mathbf{RHom}(\mathcal{A}, \mathcal{A}) & \longrightarrow & \Pi_{i>0} \mathbf{RHom} \left((\mathcal{A})^{\leq i}, \mathcal{A} \right) & \longrightarrow & \Pi_{i>0} \mathbf{RHom} \left((\mathcal{A})^{\leq i}, \mathcal{A} \right) \\ & & \downarrow \cong \text{can} & & \downarrow \cong \text{can} \\ \mathbf{RHom}(\mathcal{A}, \mathcal{B}) & \longrightarrow & \Pi_{i>0} \mathbf{RHom} \left((\mathcal{A})^{\leq i}, \mathcal{B} \right) & \longrightarrow & \Pi_{i>0} \mathbf{RHom} \left((\mathcal{A})^{\leq i}, \mathcal{B} \right). \end{array}$$

By the axioms defining a triangulated category, there exists an isomorphism $\mathbf{RHom}(\mathcal{A}, \mathcal{A}) \xrightarrow{\sim} \mathbf{RHom}(\mathcal{A}, \mathcal{B})$ making the diagram commutative; let $\mu \in \mathbf{Hom}(\mathcal{A}, \mathcal{B})$ be the image of the identity in $\mathbf{Hom}(\mathcal{A}, \mathcal{A})$ given by this isomorphism. Any morphism $\mathcal{L} \rightarrow \mathcal{A}$, with \mathcal{L} a perfect complex, factors through $\coprod_{i>0} (\mathcal{A})^{\leq i} \rightarrow \mathcal{A}$. Thus, by the commutativity of the diagram above, one sees that $\text{can}_{\mathcal{L}} : \mathbf{RHom}_{\tilde{X}}(\mathcal{L}, \mathcal{A}) \xrightarrow{\cong} \mathbf{RHom}_{\tilde{X}}(\mathcal{L}, \mathcal{B})$ coincides with μ_* . Since $\mathbf{D}(\mathbf{QCoh}(\tilde{X}))$ is compactly generated by the proof of Theorem A.2, this proves that $\mu : (\iota_X)_* \bar{f}^! \mathcal{F} \rightarrow f^! (\iota_Y)_* \mathcal{F}$ is an isomorphism. \square

We will need some notation for what follows. Given infinitesimal deformations \tilde{X} and \tilde{Y} , let $\tilde{X} \times \tilde{Y}$ denote the infinitesimal deformation $(X \times Y, \mathcal{A}_{X \times Y})$, where the restriction of the structure sheaf $\mathcal{A}_{X \times Y}$ on open sets of the form $U \times V \subset X \times Y$, with U an open affine in X and V an open affine in Y , is $(\mathcal{A}_X(U) \otimes_{R_1} \mathcal{A}_Y(V))^\sim$ (observe that $\mathcal{A}_{X \times Y}$ satisfies (**)) by Lemma 2.1.8 in [4]). Suppose $f : \tilde{X} \rightarrow \tilde{Y}$ is a flat morphism; write $\Gamma_{\iota_Y} : Y \rightarrow Y \times \tilde{Y}$ for the graph of ι_Y , $\Gamma_f : \tilde{X} \rightarrow \tilde{Y} \times \tilde{X}$ for the graph of f , and $\delta : X \rightarrow Y \times \tilde{X}$ for the morphism given by the pair (\bar{f}, ι_X) . Define $\mathcal{K}_{\iota_Y} := (\Gamma_{\iota_Y})_* \mathcal{O}_Y$, $\mathcal{K}_f := (\Gamma_f)_* \mathcal{A}_X$, and $\mathcal{K} := \delta_* \mathcal{O}_X$. Then one has isomorphisms of functors between bounded above derived categories of quasi-coherent sheaves :

$$\Phi_{\mathcal{K}_{\iota_Y}} \cong \mathbf{L}\iota_Y^*, \quad \Phi_{\mathcal{K}_f} \cong \mathbf{R}f_*, \quad \Phi_{\mathcal{K}} \cong \mathbf{R}\bar{f}_* \mathbf{L}\iota_X^*.$$

Note that in defining the Fourier–Mukai functors (see [7]) $\Phi_{\mathcal{K}_{\iota_Y}}$, $\Phi_{\mathcal{K}_f}$ and $\Phi_{\mathcal{K}}$, we have used the *bimodule* structure of the kernels for the respective structure sheaves. Consider this Cartesian diagram of infinitesimal deformations:

$$\begin{array}{ccc} X & \xrightarrow{\iota_X} & \tilde{X} \\ \bar{f} \downarrow & & \downarrow f \\ Y & \xrightarrow{\iota_Y} & \tilde{Y}. \end{array}$$

The following observation about this diagram will be used later: there is an isomorphism of functors

$$(A.1) \quad f^*(\iota_Y)_* \cong (\iota_X)_* \bar{f}^*.$$

Lemma A.5. (Kuznetsov) *There is an isomorphism $\mathcal{H}^0(\mathcal{K}_f \circ \mathcal{K}_{\iota_Y}) \cong \mathcal{K}$, inducing a morphism of functors $\mathbf{L}\iota_Y^* \mathbf{R}f_* \cong \Phi_{\mathcal{K}_{\iota_Y}} \circ \Phi_{\mathcal{K}_f} \rightarrow \Phi_{\mathcal{K}} \cong \mathbf{R}\bar{f}_* \mathbf{L}\iota_X^*$. This morphism is an isomorphism if and only if $\mathcal{K}_f \circ \mathcal{K}_{\iota_Y} \cong \mathcal{K}$, which in turn holds true if and only if $f^*(\iota_Y)_* \cong (\iota_X)_* \bar{f}^*$.*

Proof. The first statement follows essentially from the proof of Lemma 2.17 of [3]. The reader should note that while our definition of Fourier–Mukai transform is different from Kuznetsov’s, the proof still works because of the isomorphism (A.1).

For the proof of the second statement, we would like to invoke Proposition 2.15 of [3], namely, that an isomorphism of kernel functors induces an isomorphism of kernels. This result, however, depends on the existence of a perfect spanning class (see Definition 2.9 in [3]), which is false for a general infinitesimal deformation. Nevertheless, the category $D^-(\mathbf{QCoh}(\tilde{X}))$ certainly does admit a “flat spanning class”: $\{\mathcal{A}_x : x \in X\}$, where \mathcal{A}_x is the stalk of \mathcal{A}_X at x , seen as a quasi-coherent sheaf over \tilde{X} . Indeed, flatness follows from Proposition 2.1.16 (ii) of [4], whereas the spanning property, that is $H^\bullet(X, \mathcal{G} \otimes_{\mathcal{A}_X} \mathcal{A}_x) = 0$ for all $x \in X$ implies $\mathcal{G} = 0$, can be proven by the same argument as in Lemma 2.13 of [3], using flatness and the crucial observation that $H^{i>0}(M_x) = 0$ for any quasi-coherent \mathcal{A}_X -module M . This is enough to prove the analogue of Lemma 2.15 for bounded above derived categories of quasi-coherent sheaves, an easy exercise left to the reader.

The last statement is obtained simply by interpreting the kernels in the lemma as kernels of Fourier–Mukai functors going in the opposite direction. \square

In view of observation (A.1), we in fact have $\mathbf{L}\iota_Y^* \mathbf{R}f_* \cong \mathbf{R}\bar{f}_* \mathbf{L}\iota_X^*$, thus completing the proof of Lemma A.3.

Theorem A.6. *Let $f : \tilde{X} \rightarrow \tilde{Y}$ be a flat morphism, with \bar{f} smooth. Then $f^!$ commutes with coproducts and is given by the formula:*

$$f^! \mathcal{F} = f^* \mathcal{F} \otimes_{\mathcal{A}_X} f^! \mathcal{A}_Y.$$

Proof. First note that the functors $\mathbf{L}\iota_Y^*$, $\bar{f}^!$ and $(\iota_X)_*$ commute with coproducts. Then, using the second exact triangle in the proof of Proposition A.4, the first part of the result follows. For the second part, we appeal to the proof of Theorem 5.4 of [5], making use of the fact that $f^!(\mathcal{A}_Y) \in D^b(\mathbf{Coh}(\tilde{X}))$ by Proposition A.4, so that the necessary tensor products appearing there are defined (the point is that tensor products need not be defined for the unbounded derived category in the noncommutative setting). \square

Corollary A.7. *Let $\Phi_{\mathcal{E}} : D^b(\mathbf{QCoh}(\tilde{X})) \rightarrow D^b(\mathbf{QCoh}(\tilde{Y}))$ be an integral transform, with \mathcal{E} a perfect object. Then its right adjoint is an integral functor with kernel $p_{\tilde{Y}}^1(\mathcal{A}_X) \otimes \mathcal{E}^\vee$, where $p_{\tilde{Y}} : \tilde{X} \times \tilde{Y} \rightarrow \tilde{Y}$ is the second projection. In particular, if X and Y are smooth and projective, the inverse of a Fourier–Mukai equivalence between $D^b(\mathbf{Coh}(\tilde{X}))$ and $D^b(\mathbf{Coh}(\tilde{Y}))$ is a Fourier–Mukai functor.*

Denote by $\mathcal{Z}(\mathcal{A}_X)$ the center of \mathcal{A}_X , and for $\tau \in H^2(\tilde{X}, \mathcal{Z}(\mathcal{A}_X)^\times)$, let $\mathbf{QCoh}(\tilde{X}, \tau)$ and $\mathbf{Coh}(\tilde{X}, \tau)$ be the categories of twisted quasi-coherent and, respectively, twisted coherent sheaves on X (see Definition 4.1, [7]).

Corollary A.8. *Let $f : \tilde{X} \rightarrow \tilde{Y}$ be a flat morphism, with \bar{f} smooth and projective; let $\tau \in H^2(\tilde{Y}, \mathcal{Z}(\mathcal{A}_Y)^\times)$. Then, $\mathbf{R}f_* : D(\mathbf{QCoh}(\tilde{X}, f^* \tau)) \rightarrow D(\mathbf{QCoh}(\tilde{Y}, \tau))$ admits a right adjoint $f^!$, with $f^!(\mathcal{F}) = f^*(\mathcal{F}) \otimes_{\mathcal{A}_X} f^!(\mathcal{A}_Y)$, where $f^!$ in the term on the right is the untwisted right adjoint of Theorem A.2. Moreover, $f^!$ restricts to $f^! : D_{\text{perf}}(\tilde{Y}, \tau) \rightarrow D^b(\mathbf{Coh}(\tilde{X}), f^* \tau)$.*

The analogue of Corollary A.7 also holds in this setting.

Proof. This is standard given the previous results in the untwisted setting (see, for example, Theorem 2.4.1 of [2]). \square

Remark A.9. Let us define *deformations of order n* as locally ringed spaces of the form $(X, \mathcal{A}_{X,n})$, where X is a separated, noetherian scheme over \mathbb{C} , and $\mathcal{A}_{X,n}$ is a sheaf of local flat $R_n := \mathbb{C}[\epsilon]/\mathbb{C}[\epsilon^{n+1}]$ -algebras with a fixed isomorphism $\mathcal{A}_X \otimes_{R_n} \mathbb{C} \cong \mathcal{O}_X$, satisfying the following recursive conditions:

- (*) $0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{A}_{X,n} \rightarrow \mathcal{A}_{X,n-1} \rightarrow 0$ is a central extension (with $\mathcal{A}_{X,n-1}$ a deformation of order $n-1$),
- (**') for any local section $f \in \mathcal{O}_X(U)$ over an affine open set U , there is a lift $\hat{f} \in \mathcal{A}_{X,n}(U)$ such that the multiplicative set $\{\hat{f}^n : n \geq 0\}$ respects the left and right Ore localization conditions.

Then, it is not hard to see by dévissage, using the exact triangles arising from the exact sequence $0 \rightarrow \mathbb{C} \rightarrow R_{i+1} \rightarrow R_i \rightarrow 0$, as in the proof of Proposition A.4, that the results of this appendix carry over to order n . The usefulness of this generalization may be limited, however. Indeed, while every infinitesimal deformation of the abelian category $\mathbf{Coh}(\mathcal{O}_X)$ for a smooth and projective variety X admits a description as the twisted category $\mathbf{Coh}(\tilde{X}, \tau)$ of an infinitesimal deformation $(\tilde{X}, \mathcal{A}_X)$ in our sense, it is not clear to us what form a general higher order deformation takes.

REFERENCES

- [1] M. Bökstedt, A. Neeman, *Homotopy limits in triangulated categories*, *Compositio Math.* **86** (1993), 209–234.
- [2] A. Căldăraru, *Derived categories of twisted sheaves on Calabi-Yau manifolds*, Cornell Univ. Ph.D. Thesis (2000), <http://www.math.wisc.edu/~andreic/>.
- [3] A. Kuznetsov, *Hyperplane sections and derived categories*, *Izv. Math.* **70** (2006), 447–547.
- [4] J. C. McConnell, J. C. Robson, *Noncommutative Noetherian Rings*, Graduate Studies in Mathematics Vol. 30, American Mathematical Society, Providence, RI, 2001.
- [5] A. Neeman, *The Grothendieck Duality Theorem via Bousfield's techniques and Brown Representability*, *J. Amer. Math. Soc.* **1** (1996), 205–236.
- [6] N. Spaltenstein, *Resolution of unbounded complexes*, *Compositio Math.* **65** (1988), 121–154.
- [7] Y. Toda, *Deformations and Fourier–Mukai transforms*, *J. Diff. Geom.* **81** (2009), 197–224.

E.M.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, 155 SOUTH 1400 EAST, SALT LAKE CITY, UT 84112, USA

E-mail address: macri@math.utah.edu

S.M.: DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS–AMHERST, 710 N. PLEASANT STREET, AMHERST, MA 010003, USA

E-mail address: mehrotra@math.umass.edu

P.S.: DIPARTIMENTO DI MATEMATICA “F. ENRIQUES”, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA CESARE SALDINI 50, 20133 MILANO, ITALY

E-mail address: paolo.stellari@unimi.it

URL: <https://sites.unimi.it/stellari>